

УДК 550.47:581.526.3(476)

ПРОСТРАНСТВЕННАЯ ИЗМЕНЧИВОСТЬ СОДЕРЖАНИЯ ТЯЖЕЛЫХ МЕТАЛЛОВ В МАКРОФИТАХ ОЗЕР БЕЛАРУСИ

Н. В. ЖУКОВСКАЯ¹⁾, Н. В. КОВАЛЬЧИК¹⁾, Б. П. ВЛАСОВ¹⁾, Т. МОЛЕНДА²⁾

¹⁾Белорусский государственный университет, пр. Независимости, 4, 220030, г. Минск, Беларусь

²⁾Силезский университет в Катовицах, ул. Бендзиньска, 60, 41-200, г. Сосновец, Польша

Проанализирована пространственная дифференциация концентрации тяжелых металлов (Ti, V, Cr, Mn, Ni, Cu, Pb) в высшей водной растительности озер Беларуси. Данные о содержании элементов в макрофитах Беларуси сопоставимы с показателями, полученными в Польше и других соседних странах. Озерным растениям свойственна избирательность в накоплении металлов, различающаяся для экологических групп погруженных, надводных растений и макрофитов с плавающими листьями. Установлены статистически значимые отличия между бассейнами рек по содержанию Mn, Cu, Pb в погруженных макрофитах озер. Наибольшими концентрациями Mn в тканях макрофитов всех экологических групп характеризуются бассейны рек Днепр, Припять, Западный Буг, наименьшими – р. Березины. Наименьшее количество Pb обнаружено в водных растениях озер бассейнов рек Западный Буг и Днепр. Водные растения озер различной степени трофности отличаются на статистически значимом уровне по содержанию Mn и Cu, наименьшие концентрации Mn в тканях макрофитов характерны для мезотрофных озер с признаками олиготрофии. С помощью пространственного кластерного анализа выделены восемь групп озер по содержанию Mn, Cu и Pb в высшей водной растительности. Группа озер Браславского района характеризуется повышенным содержанием Mn и Cu в тканях макрофитов, озера бассейна р. Вилии отличаются наименьшей среди других групп средней концентрацией Cu и Pb в водных растениях, озера северной части Беларуси выделяются повышенным содержанием Cu и Pb в тканях растений.

Ключевые слова: тяжелые металлы; высшая водная растительность; экологические группы; пространственная изменчивость; мониторинг; трофический статус озер.

Образец цитирования:

Жуковская НВ, Ковальчик НВ, Власов БП, Моленда Т. Пространственная изменчивость содержания тяжелых металлов в макрофитах озер Беларуси. *Журнал Белорусского государственного университета. География. Геология.* 2019; 2:40–51 (на англ.).
<https://doi.org/10.33581/2521-6740-2019-2-40-51>

For citation:

Zhukovskaya NV, Kavalchik NV, Vlasov BP, Molenda T. Spatial variations of trace elements contents in macrophytes within Belarusian lakes. *Journal of the Belarusian State University. Geography and Geology.* 2019;2:40–51.
<https://doi.org/10.33581/2521-6740-2019-2-40-51>

Авторы:

Наталья Викторовна Жуковская – кандидат географических наук, доцент; доцент кафедры почвоведения и географических информационных систем факультета географии и геоинформатики.

Надежда Владимировна Ковальчик – кандидат географических наук, доцент; доцент кафедры почвоведения и географических информационных систем факультета географии и геоинформатики.

Борис Павлович Власов – доктор географических наук, профессор; заведующий научно-исследовательской лабораторией озераведения факультета географии и геоинформатики.

Тадеуш Моленда – доктор географических наук; доцент кафедры физической географии факультета наук о Земле.

Authors:

Natalia V. Zhukovskaya, PhD (geography), docent; associate professor at the department of soil science and geographic information systems, faculty of geography and geoinformatics.
natazhuk@gmail.com

Nadzeja V. Kavalchik, PhD (geography), docent; associate professor at the department of soil science and geographic information systems, faculty of geography and geoinformatics.
kovalchiknv@gmail.com

Boris P. Vlasov, doctor of science (geography), full professor; head of the laboratory of lakes research, faculty of geography and geoinformatics.
vlasov@bsu.by

Tadeusz Molenda, doctor of science (geography); associate professor at the department of physical geography, faculty of earth science.
tedimolenda@interia.pl

SPATIAL VARIATIONS OF TRACE ELEMENTS CONTENTS IN MACROPHYTES WITHIN BELARUSIAN LAKES

N. V. ZHUKOVSKAYA^a, N. V. KAVALCHYK^a, B. P. VLASOV^a, T. MOLEND^b

^aBelarusian State University, 4 Niezaliežnasci Avenue, Minsk 220030, Belarus

^bUniversity of Silesia in Katowice, 60 Będzińska Street, Sosnowiec 41-200, Poland

Corresponding author: N. V. Zhukovskaya (natazhuk@gmail.com)

The spatial variation of trace elements contents of heavy metals (Ti, V, Cr, Mn, Ni, Cu, Pb) in higher aquatic plants of Belarusian lakes has been analyzed according to monitoring data. Natural variability limits of metals contents in aquatic plants of lakes within principal rivers basins of Belarus have been determined. The data on trace elements contents in macrophytes of Belarus are comparable with the data obtained in Poland and other neighbouring countries. Aquatic plants in lakes are characterized by selective metals accumulation, which is different for ecological groups of submersed, emergent and macrophytes with floating leaves. Statistically significant differences between river basins in the contents of Mn, Cu and Pb have been established in the submersed macrophytes. For all ecological groups, the highest concentrations of Mn in lakes macrophytes tissues are observed in the basins of the Dniapro, Prypiač, Zachodni Buh, and the lowest – the Biarezina. The lakes of the Zachodni Buh and Dniapro basins are distinguished by lower Pb concentrations in aquatic plants. Significant differences in the content of Mn and Cu in the plant tissues within different trophicity lakes have been established. The lowest concentrations of Mn are observed in mesotrophic lakes with signs of oligotrophy. Group analysis of lakes based on the content of Mn, Cu, Pb in macrophyte tissues and spatial constraints has been performed. Among the allocated groups are the lakes of the Braslav district characterized by elevated content of Mn and Cu; lakes of the Vilija basin, distinguished by the lowest average levels of Cu and Pb; lakes of the northern part of Belarus, notable for increased contents of Cu and Pb.

Keywords: trace elements; higher aquatic plants; ecological groups; spatial variability; monitoring; lakes trophic level.

Introduction

Aquatic macrophytes may accumulate chemical elements in concentrations exceeding their content in the environment. The concentrations of chemical elements in aquatic plants can be more than 100 000 times higher than in the associated water [1]. This accumulation activity determines their use in the system of monitoring higher aquatic plants and monitoring the state of water bodies in the National Environment Monitoring System of the Republic of Belarus. A major limit to use macrophytes as bioindicators is the variability of elements in the organisms. The most important causes of the variability are a seasonal cycle, the biological state of organisms (species, size, age, and probably growth rate) and physical environment [2]. Chemical composition of water and bottom sediments is the result of climatic, hydrological, mechanical, physical, chemical, biological and others processes occurring both on the catchment area and in the reservoir itself. The aim of this paper is to identify the spatial characteristics of trace elements (Ti, V, Cr, Mn, Ni, Cu, Pb) accumulated by aquatic plants in Belarusian lakes.

Materials and methods

In the present study we have used data from the National Environment Monitoring System. The accumulation of heavy metals by higher aquatic plants within Belarus reservoirs and lakes from 2000 to 2015 has been analyzed. A monitoring network on lakes includes 46 key sites (fig. 1). The representativeness of the network is ensured by the monitoring objects distribution in accordance with the lake percentage and physiographic zoning of Belarus. The lakes differ in the hollows origin, morphometry, and degree of overgrowing.

Bottom sediments in macrophytes growing locations are mainly represented by sand and silted sand in littoral zones and deeper sandy silts. The samples were formed from macrophytes tissues of various ecological groups and have been considered in this study: emergent (*Acorus calamus* L., *Phragmites australis* (Cav.) Trin. ex Steud., *Carex rostrata* Stokes, *Carex lasiocarpa* Ehrh. and *Carex acuta* L., *Sparganium erectum* L., *Sparganium natans* L., *Glyceria maxima* (O. Hartm) Holmb, *Glyceria fluitans* (L.) R. Br., *Cladium mariscus* (L.) Pohl, *Schoenoplectus lacustris* (L.), *Scirpus sylvaticus* L., *Typha angustifolia* L., *Typha latifolia* L., *Sagittaria sagittifolia* L., *Butomus umbellatus* L., *Equisetum fluviatile* L., *Zizania palustris* L., *Zizania latifolia* (Griseb.) Stapf, *Alisma plantago-aquatica* L.); submersed (*Hydrilla verticillata* (L. fil.) Royle, *Potamogeton lucens* L., *Potamogeton perfoliatus* L., *Potamogeton trichoides* Cham. et Schlecht., *Potamogeton praelongus*




Fig. 1. Macrophyte sampling locations

Wulf., *Potamogeton crispus* L., *Potamogeton acutifolius* Link, *Potamogeton compressus* L., *Potamogeton obtusifolius* Mert. et Koch, *Potamogeton nodosus* Poir., *Potamogeton friesii* Rupr., *Myriophyllum spicatum* L., *Myriophyllum verticillatum* L., *Elodea canadensis* Michx., *Ceratophyllum submersum* L., *Ceratophyllum demersum* L., *Stratiotes albidus* L., *Isoetes lacustris* L., *Littorella uniflora* (L.) Aschers); floating-leaved (*Persicaria amphibian* (L.), *Nuphar lutea* (L.) Smith., *Nymphaea candida* J. et C. Presl., *Potamogeton natans* L.). Charophyta, which are often referred to as higher aquatic vegetation.

The time of sampling corresponds to maximum biomass development and chemicals accumulation by plants during the flowering period (July – August).

Analytical determination of trace elements (Ti, V, Cr, Mn, Ni, Cu, Pb) is performed by emission spectral analysis methods described by Zyrin, Obukhov [3].

For the analysis and visualization of spatial variability of trace elements contents in lake macrophytes, ArcGIS 10.3 geographic database has been designed and developed. Vectors datasets have been created based on OpenStreetMap data¹. Point feature class «lakes_monitoring» contains characteristics of monitoring points. The relationship class has been created to associate a table with data on trace elements contents in macrophytes and the «lakes_monitoring» feature class. Watersheds of principal rivers within Belarus have been identified based on SRTM digital relief model² using ArcGIS Hydrology tools.

Statistical data analysis includes the variable distribution assessment with the help of histograms, normal quantile – quantile (Q–Q) plots and fitting criterions (the Kolmogorov – Smirnov test, the Shapiro – Wilk test), as well as data transformation, calculating descriptive statistics, the Kruskal – Wallis test which is the most effective for samples with a markedly different number of observations. For the lognormal (ln) distribution, the geometric mean (g) is taken as the average. In the case of truncated samples, with frequency of trace elements occurrence more than 50 % the median is used as the average. The level of significance is set at $p \leq 0.01$ (two-tailed).

¹Download OpenStreetMap data for this region: Europe [Electronic resource]. URL: <http://download.geofabrik.de/europe.html> (date of access: 11.01.2019).

²SRTM data // CGIAR – Consortium for Spatial Information (CGIAR-CSI) [Electronic resource]. URL: <http://srtm.csi.cgiar.org/srtmdata/> (date of access: 11.01.2019).

Cluster analysis of the examined lakes on the base of trace elements contents in macrophyte tissues and spatial limitation has been performed using the ArcGIS Grouping Analysis tool (uses a K-means algorithm).

Distribution parameters of trace elements in lakes macrophytes are compared in terms of principal Belarusian river basins by ecological groups of aquatic plants: submersed, floating-leaved, emergent.

Results and discussion

Geochemical features of aquatic plants are largely determined by their membership in ecological groups, among which systematic differences can appear [4]. Data analysis has shown that the average elemental contents are different in aquatic plants groups. The highest concentrations of all analyzed elements are characteristic of submersed macrophytes (table 1).

Such a high accumulation capability of submersed macrophytes in relation to trace elements was noted by many authors [5; 6]. Submersed plant leaves have a very thin cuticle. The leaves of submersed plants are therefore very good at taking up metals directly from the water [7].

Table 1

**Metals contents in macrophytes
of various ecological groups within reservoirs**

Element (frequency, %)	Submersed, mg/kg (n = 305)	Floating-leaved, mg/kg (n = 132)	Emergent, mg/kg (n = 320)
Ti (51–67)	4.3 (n. d. – 973)	1.28 (n. d. – 305)	1.18 (n. d. – 184)
V (30–39)	5.37 (n. d. – 141)	2.01 (n. d. – 19.0)	2.34 (n. d. – 19.3)
Cr (23–42)	4.32 (n. d. – 94.2)	0.58 (n. d. – 7.32)	0.85 (n. d. – 14.8)
Mn (100)	296 (0.60–9158)	101.4 (0.23–3957)	23.1 (0.03–3259)
Ni (23–40)	1.18 (n. d. – 23.6)	0.094 (n. d. – 0.87)	0.39 (n. d. – 23.3)
Cu (100)	2.25 (0.07–235.6)	0.32 (0.07–12.2)	1.37 (0.05–76.1)
Pb (100)	2.93 (0.08–471.2)	0.39 (0.04–18.3)	0.42 (0.03–93.1)

Note. N. d. means not detected.

Ti, V, Mn and Pb concentrations in all examined aquatic plants are within the range described by Markert [8]. Cu and Ni abundance is lower than the average content in a plant. The data on trace elements contents in macrophytes of Belarus are mainly comparable with the data obtained in Poland and other neighboring countries (table 2).

Table 2

**Elemental composition of aquatic macrophytes obtained
in this study compared with data from [5; 8–14], mg/kg dry wt**

Indicator	Ti	V	Cr	Mn	Ni	Cu	Pb
Average contents in plant [8]	0.02–56.0	0.001–10.0	0.02–1.0	1–700	0.4–4.0	5–20	0.1–5.0
Normal level in plant leaves [9]	–	0.2–1.5	0.1–1.5	30–300	0.1–1.5	5–30	5–10
Poland							
Slup Dam Reservoir [10]							
<i>Potamogeton crispus</i> L.	–	–	–	–	5.9	17.5	5.4
<i>Ceratophyllum demersum</i> L.	–	–	–	–	6.9	9.7	5.6
<i>Phragmites communis</i> Trin.	–	–	–	–	5.7–8.0	5.4–12.3	2.6–9.9
Dominickie Lake [11]							
<i>Ceratophyllum demersum</i> L.	–	–	1.4	304	1.28	2.5	16.9
West Poland lakes [12]							
<i>Nymphaea alba</i>	–	–	1.2–4.7	88–247	1.1–1.2	1.7–4.8	8.7–4.8
<i>Nuphar luteum</i>	–	–	1.4–3.1	253–345	1.4–1.6	1.8–2.2	10.5–11.9
Lake Symsar [12]							
<i>Myriophyllum spicatum</i> L.	–	–	3.9	1190	24	4.2	6.0
<i>Nymphaea alba</i> L.	–	–	1.0	510	3.0	0.9	Less 1
<i>Typha angustifolia</i> L.	–	–	3.6	540	23	3.3	15

Ending table 2

Indicator	Ti	V	Cr	Mn	Ni	Cu	Pb
The Russian Federation							
Bugach Lake [13] Macrophytes	297	1.71	8.66	407	7.30	50.5	0.57
Ivan'kovskoe Reservoir [5]							
<i>Potamogeton perfoliatus</i> L.	–	2.6	5.0	1240	7.1	8.8	1.4
<i>Ceratophyllum demersum</i> L.	–	0.44	11.8	5260	14	55.3	4.5
<i>Myriophyllum spicatum</i> L.	–	Less 0.45	3.0	2898	1.8	14.5	1.0
Ukraine							
Zaporozhian Reservoir [14]							
<i>Potamogeton perfoliatus</i> L.	–	–	–	960–1830	6.2–19.0	6–18	3.0–6.1
<i>Phragmites australis</i>	–	–	–	29–112	1.7–4.6	3.1–6.2	0.7–1.3
Belarus							
Lakes and reservoirs							
<i>Potamogeton perfoliatus</i> L.	2.7	3.4	1.8	186	0.41	1.4	1.8
<i>Potamogeton crispus</i> L.	4.0	0.66	1.7	225	0.99	1.8	2.77
<i>Myriophyllum spicatum</i> L.	4.0	5.6	4.7	269	1.15	2.8	4.4
<i>Ceratophyllum demersum</i> L.	1.3	5.4	6.2	702	1.2	3.3	4.3
<i>Nymphaea alba</i> L.	3.0	2.1	0.25	69	0.11	0.28	0.18
<i>Nuphar luteum</i>	0.14	1.6	0.34	79	0.07	0.25	0.18
<i>Phragmites australis</i>	0.05	2.1	0.35	11.8	0.09	2.0	0.49
<i>Typha angustifolia</i> L.	1.47	0.35	0.20	110.0	0.02	1.2	0.17

Non-parametric dispersion analysis of Kruskal – Wallis has been used to compare Mn, Cu and Pb contents (elements with 100 % occurrence) in lake macrophytes. The results of the test demonstrate statistically significant differences between river basins in Mn ($H = 26.7$, $df = 6$, $p < 0.001$), Cu ($H = 23.3$, $df = 6$, $p < 0.001$) and Pb content ($H = 31.8$, $df = 6$, $p < 0.001$) in submersed macrophytes; Mn ($H = 32.7$, $df = 6$, $p < 0.001$) and Pb ($H = 17.0$, $df = 6$, $p < 0.009$) content in emergent macrophytes; Mn ($H = 31.2$, $df = 6$, $p < 0.001$) content in submersed macrophytes with floating leaves.

The highest spatial variations is noted in taking up of Mn by macrophytes (fig. 2). The average Mn content in submerged macrophytes within different catchments ranges from 1736 (the Dniapro River Basin) to 208 mg/kg (the Biarezina River Basin).

A similar pattern of spatial variability for examined trace element contents in lakes macrophytes for all ecological groups has been revealed (fig. 2 and 3). The highest concentrations of Mn in lakes macrophytes tissues are observed in the basins of the Dniapro, Prypiac, Zachodni Buh, and the lowest – in the Biarezina. The lakes of the Zachodni Buh and Dniapro basins are distinguished by the lowest Pb contents in aquatic plants. The average rank variation of Mn, Cu and Pb contents in submersed and emergent macrophytes in river basins is presented in fig. 4 and 5. The analyzed elements are distinguished by the spatial variability nature. Spatial variation of Pb and Cu contents in submersed as well as emergent macrophytes at principal river basins level is similar in nature.

Metals concentration levels in macrophytes depends on a large number of various factors. The chemical composition of aquatic plants reflects the geochemical features of lakes bottom sediments, rocks and soils within catchment areas and it depends on lake morphometrical characteristics and production processes, on tehnogenesis products flow into reservoirs from local sources.

In general, trace elements contents in rocks and soils of Belarus are lower than the average world values. So, Belarusian soils are depleted of Ti, Mn, V, Ni, Cu about 3 times, Cr – 2 times, the Pb content is comparable to Clarke values [15]. In lake sediments elements accumulate in lower concentrations than in soils, with the exception of Mn and Pb. Glacial complexes of the poozersky and pripyat ages [16], as well as poozersko-holocene alluvial, lake-alluvial and aeolian accumulations, holocene alluvial and bog formations prevail among the covering deposits. In some areas, due to a variegated lithological composition, the covering deposits can be characterized by elevated trace elements concentrations [17].

The territory within the Zachodniaya Dzvina River Basin is characterized by the highest lake density. In conditions of a significant vertical dissection of relief the lake hollows have a deep (up to 30–50 m) erosional incision, and the catchments have been developed and plowed up. A lot of allochthonic terrigenous material enters the reservoirs. Silicate type of sedimentation is typical for lakes [18]. The soils within the Zachodniaya Dzvina River Basin are distinguished by the highest concentrations of Cu, Ni, Cr, to a lesser extent Ti, the lowest – Zr [19].

Within the basins of the Viliya and the Nioman as well as the Biarezina (the right tributary of the Dniapro) both low-flowing lakes with organo-silicate, mixed and carbonate sediments, and biogenic lithogenesis lakes with an increased amount of organic matter prevail. The soils of the Nioman River Basin are characterized by a high content of Mn, low – Ni, Co and Cr.

The territories of the Dniapro and Sož river basins are distinguished by low lakes density, well-defined surface runoff and the development of suffusion processes. A reduced content of trace elements except Ni has been noted, the accumulation of which is associated with the minerals aleuritic fraction that predominates in the composition of covering deposits [20]. The low forest cover within the Dniapro Basin (22 %) indicates an increase in the intensity of the input of biogenic and terrigenous material into water bodies.

In the Zachodni Buh Basin, there is a high level of forest cover (60 %), water-glacial plains dominate with sandy soils which are poor of trace elements. Within the Prypiač River Basin, extensive plains of river terraces and sandurs with swampy lowerings and flat lake hollows are widespread. Lake density is low, silicate-organic, organo-silicate, mixed and carbonate sediments form in weak-flow lakes. The geochemical specificity of soils in this basin reveals Fe accumulation, low contents of Mg and trace elements (Ti, Mn, Ni, Co, Cr) compared with the average values in Belarus [19].

The modern formation of lake sediments on the territory of Belarus is influenced by technogenesis. There is increasing the flow of terrigenous material (mainly clayey) from watersheds (with which Ti, Mn, Zn, Cr, V, Pb are associated) as a result of the agricultural and reclamation use of the land. The geochemical mobility of Mn, Cu, Zn is increasing as a result of organo-mineral interaction [18]. Some abnormal concentrations of Pb, Ni, Cu, Cr in lake sediments are associated with the impact of local technogenic sources [21].

Direct inheritance of the chemical composition of bottom sediments from elements content in soils and rocks within catchment areas has not been revealed. Farhat and Aly [22] suggested that the organic matter was more important in controlling heavy metals distribution. Statistically significant relationships between analyzed trace elements contents in macrophytes tissues and bottom sediments has been determined only for Mn.




Fig. 2. Relative trace element abundances in submersed macrophytes of Belarusian lakes within principal river basins




Fig. 3. Relative trace element abundances in emerged (a) and submersed with floating leaves (b) macrophytes of Belarusian lakes within principal river basins




Fig. 4. Average rank variation of Mn, Cu, Pb contents in submersed lake macrophytes by principal river basins




Fig. 5. Average rank variation of Mn, Cu, Pb contents in emerged lake macrophytes by principal river basins

A fairly complete picture of the ecological conditions for the growth of macrophytes gives the degree of trophic water bodies. The effect of trophic status on trace elements contents in aquatic plants was estimated using the example of submersed macrophytes (table 3). Statistically significant differences in Mn and Cu contents in the tissues of plants growing in lakes of varying trophicity have been established (H variates from 16.2 to 16.9, $p < 0.01$). The lowest concentration of Mn has been observed in plant tissues of mesotrophic lakes with signs of oligotrophy. This dependence is also typical of plants in other ecological groups (fig. 6).

It should be noted that within the Zachodni Buh and Dniapro basins standing out for high Mn content in aquatic plants, eutrophic lakes make up 100 % of the sample (fig. 7).

Cluster analysis of monitoring points based on Mn, Cu and Pb contents in macrophyte tissues and spatial constraints has been performed. Eight clusters have been allocated (fig. 8).

Table 3

Trace elements contents in submersed plants within lakes of different trophic status, mg/kg

Trophic status	Ti	V	Cr	Mn	Ni	Cu	Pb
Mesotrophic lakes with signs of oligotrophy ($n = 64$)	3.87	9.24	7.37	125	2.04	2.84	3.1
Mesotrophic ($n = 87$)	13.6	6.38	4.45	363	1.23	3.18	4.2
Eutrophic ($n = 146$)	1.65	3.16	3.12	464	0.76	1.82	2.39
Dystrophic ($n = 8$)	0.14	2.30	–	237	–	0.35	2.04




Fig. 6. Mn average content in macrophytes within lakes of different trophic status, mg/kg




Fig. 7. Percentage of different trophic status lakes, %

The first group consists of the Viliya Basin lakes, which are distinguished by lower average levels of the elements (Mn, Cu, Pb). The second group includes examined lakes of the Braslav district characterized by elevated content of Mn and Cu (fig. 9). Bredna Lake (maximum Pb content), Lieskavičy Lake (maximum Mn content) and Sviciaž Lake (elevated Pb and Cu contents) form separate groups. Lakes of the northern part of Belarus, which are notable for elevated contents of Cu and Pb are combined into the sixth group. The lakes of the Zachodni Buh and Prypiac basins form a separate group described by elevated Mn concentration in macrophytes. The latter cluster includes lakes within the Biarezina, Dniapro and Sož basins and the southern part of the Zachodniaja Dzvina Basin.




Fig. 8. Cluster analysis of lakes based on the content of Mn, Cu, Pb in macrophyte tissues




Fig. 9. Mn, Cu and Pb average contents (standardized values) in submersed macrophytes of identified lakes groups

Conclusion

The variation of trace elements contents in lake macrophytes on principal river basins has been investigated. Statistically significant differences between river basins in the contents of Mn, Cu and Pb in submersed macrophytes, Mn and Cu in emergent macrophytes and Mn in macrophytes with floating leaves have been established. For all ecological groups the highest concentrations of Mn in lake macrophytes tissues are observed in the basins of the Dniapro, the Prypiac and the Zachodni Buh, and the lowest – in the Biarezina. The lakes of the Zachodni Buh and Dniapro basins are distinguished by the lowest Pb contents in aquatic plants.

The lakes trophic status is an important factor determining the spatial variability of trace elements accumulation in macrophyte tissues. Within the Zachodni Buh and Dniapro basins standing out for the highest Mn content in aquatic plants, eutrophic lakes make up 100 % of the sample.

Based on Mn, Cu and Pb contents in macrophyte tissues and spatial constraints eight lakes clusters has been allocated. Among the groups are the lakes of the Braslav district characterized by elevated content of Mn and Cu; lakes of the Vilija Basin, which are distinguished by lower average levels of the elements; lakes of the northern part of Belarus, which are notable for elevated contents of Cu and Pb. Separate groups form Bredna Lake (maximum Pb content) and Sviciaz Lake (elevated Pb and Cu contents).

Cognition of spatial distribution features of trace elements contents in macrophytes tissues contributes to the improvement in the system of monitoring higher aquatic plants and monitoring the state of water bodies in general.

Библиографические ссылки

1. Albers PH, Camardese MB. Effects of acidification on metal accumulation by aquatic plants and invertebrates. 1. Constructed wetlands. *Environmental Toxicology and Chemistry*. 1993 June;12(6):959–967. DOI: 10.1002/etc.5620120602.
2. Ravera O, Cenci R, Beone GM, Dantas M, Lodigiani P. Trace element concentrations in freshwater mussels and macrophytes as related to those in their environment. *Journal of Limnology*. 2003;62(1):61–70. DOI: 10.4081/jlimnol.2003.61.
3. Зырин НГ, Обухов АИ. *Спектральный анализ почв, растений и других биологических материалов*. Москва: Издательство МГУ; 1977. 334 с.
4. Гигевич ГС, Власов БП, Вынаев ГВ. *Высшие водные растения Беларуси: эколого-биологическая характеристика, использование и охрана*. Минск: Издательский центр БГУ; 2001. 231 с.
5. Гришпанцева ЕС, Сафронова НС, Кирпичникова НВ, Федорова НП. Распределение микроэлементов в высшей водной растительности Иваньковского водохранилища. *Геоэкология. Инженерная геология. Гидрогеология. Геокриология*. 2010;3: 223–231.
6. Crowder A, Dushenko WT, Greig J, Poland JS. Metal contamination in sediments and biota of the Bay of Quinte, Lake Ontario, Canada. *Hydrobiologia*. 1989 December;188(1):337–343. DOI: 10.1007/BF00027798.
7. Prasad MNV, Greger M, Aravind P. Biogeochemical cycling of trace elements by aquatic and wetland plants. In: Prasad MNV, Sajwan KS, Naidu R, editors. *Trace elements in the environment: biogeochemistry, biotechnology and bioremediation*. Boca Raton: Taylor & Francis; 2005. p. 451–482. DOI: 10.1201/9781420032048.ch24.
8. Markert B. Presence and significance of naturally occurring chemical elements of the periodic system in the plant organism and consequences for future investigations on inorganic environmental chemistry in ecosystems. *Vegetatio*. 1992 November;103(1):1–30. DOI: 10.1007/BF00033413.
9. Kabata-Pendias A. *Trace elements in soils and plants*. 4th edition. Boca Raton: CRC Press; 2010. 548 p.
10. Senze M, Kowalska-Góralaska M, Pokorny P. Metals in chosen aquatic plants in a lowland dam reservoir. *Journal of Elementology*. 2009 January;14(1):147–156. DOI: 10.13140/2.1.1422.9126.
11. Szymanowska A, Samecka-Cymermana A, Kempers AJ. Heavy Metals in Three Lakes in West Poland. *Ecotoxicology and Environmental Safety*. 1999 May;43(1):21–29. DOI: 10.1006/eesa.1998.1747.
12. Kuriata-Potasznik A, Szymczyk S, Pilejczyk D. Effect of bottom sediments on the nutrient and metal concentration in macrophytes of river-lake systems. *Annales de Limnologie – International Journal of Limnology*. 2018;54. Article number 1. DOI: 10.1051/limn/2017028.
13. Иванова ЕА, Анищенко ОВ, Грибовская ИВ, Зиненко ГК, Назаренко НС, Немчинов ВГ и др. Содержание металлов в высших водных растениях в небольшом сибирском водохранилище. *Сибирский экологический журнал*. 2012;4:485–495.
14. Федоненко ЕВ, Филиппова ЕВ. Оценка уровня загрязнения Запорожского водохранилища тяжелыми металлами. *Ученые записки Таврического национального университета имени В. И. Вернадского. Серия: Биология. Химия*. 2008;21(2):133–138.
15. Петухова НН, Кузнецов ВА. Кларкам микроэлементов в почвенном покрове Беларуси. *Доклады Академии наук Беларуси*. 1992;36(5):461–465.
16. Кручек СА, редактор. *Стратиграфические схемы докембрийских и фанерозойских отложений Беларуси: объяснительная записка*. Минск: Белорусский научно-исследовательский геолого-разведочный институт; 2010. 282 с.
17. Матвеев АВ, Бордон ВЕ. *Геохимия четвертичных отложений Беларуси*. Минск: Беларуская навука; 2013. 191 с.
18. Жуховицкая АЛ, Генералова ВА. *Геохимия озер Белоруссии*. Минск: Наука и техника; 1991. 202 с.
19. Петухова НН. *Геохимия почв Белорусской ССР*. Минск: Наука и техника; 1987. 231 с.
20. Лукашев КИ, редактор. *Геохимические провинции покровных отложений БССР*. Минск: Наука и техника; 1969. 473 с.
21. Лукашев ОВ, Жуковская НВ, Лукашова НГ, Творонович-Севрук ДЛ, Савченко СВ. Содержание химических элементов в современных донных отложениях рек Беларуси (техногеохимический аспект). *Природопользование*. 2015;27:109–117.
22. Farhat HI, Aly W. Effect of site on sedimentological characteristics and metal pollution in two semi-enclosed embayments of great freshwater reservoir: Lake Nasser, Egypt. *Journal of African Earth Sciences*. 2018;141:194–206. DOI: 10.1016/j.jafrearsci.2018.02.012.

References

1. Albers PH, Camardese MB. Effects of acidification on metal accumulation by aquatic plants and invertebrates. 1. Constructed wetlands. *Environmental Toxicology and Chemistry*. 1993 June;12(6):959–967. DOI: 10.1002/etc.5620120602.
2. Ravera O, Cenci R, Beone GM, Dantas M, Lodigiani P. Trace element concentrations in freshwater mussels and macrophytes as related to those in their environment. *Journal of Limnology*. 2003;62(1):61–70. DOI: 10.4081/jlimnol.2003.61.
3. Zyrin NG, Obukhov AI. *Spektral'nyi analiz pochv, rastenii i drugikh biologicheskikh materialov* [Spectral analysis of soils, plants and other biological materials]. Moscow: Publishing House of the Moscow State University; 1977. 334 p. Russian.
4. Gigevich GS, Vlasov BP, Vynaev GV. *Vysshie vodnye rasteniya Belarusi: ekologo-biologicheskaya kharakteristika, ispol'zovanie i okhrana* [Higher aquatic plants of Belarus: ecological and biological characteristics, use and protection]. Minsk: Publishing Center of Belarusian State University; 2001. 231 p. Russian.
5. Grishantseva ES, Safronova NS, Kirpichnikova NV, Fedorova NP. Distribution of microelements in higher aquatic plants in the Ivan'kovskoe water reservoir. *Geojekologija. Inzhenernaja geologija. Gidrogeologija. Geokriologija*. 2010;3:223–231. Russian.
6. Crowder A, Dushenko WT, Greig J, Poland JS. Metal contamination in sediments and biota of the Bay of Quinte, Lake Ontario, Canada. *Hydrobiologia*. 1989 December;188(1):337–343. DOI: 10.1007/BF00027798.
7. Prasad MNV, Greger M, Aravind P. Biogeochemical cycling of trace elements by aquatic and wetland plants. In: Prasad MNV, Sajwan KS, Naidu R, editors. *Trace elements in the environment: biogeochemistry, biotechnology and bioremediation*. Boca Raton: Taylor & Francis; 2005. p. 451–482. DOI: 10.1201/9781420032048.ch24.
8. Markert B. Presence and significance of naturally occurring chemical elements of the periodic system in the plant organism and consequences for future investigations on inorganic environmental chemistry in ecosystems. *Vegetatio*. 1992 November;103(1):1–30. DOI: 10.1007/BF00033413.
9. Kabata-Pendias A. *Trace elements in soils and plants*. 4th edition. Boca Raton: CRC Press; 2010. 548 p.
10. Senze M, Kowalska-Góralaska M, Pokorny P. Metals in chosen aquatic plants in a lowland dam reservoir. *Journal of Elementology*. 2009 January;14(1):147–156. DOI: 10.13140/2.1.1422.9126.
11. Szymanowska A, Samecka-Cymermana A, Kempers AJ. Heavy Metals in Three Lakes in West Poland. *Ecotoxicology and Environmental Safety*. 1999 May;43(1):21–29. DOI: 10.1006/eesa.1998.1747.
12. Kuriata-Potasznik A, Szymczyk S, Pilejczyk D. Effect of bottom sediments on the nutrient and metal concentration in macrophytes of river-lake systems. *Annales de Limnologie – International Journal of Limnology*. 2018;54. Article number 1. DOI: 10.1051/limn/2017028.
13. Ivanova EA, Anishchenko OV, Gribovskaya IV, Zinenko GK, Nazarenko NS, Nemchinov VG, et al. Metal content in higher aquatic plants in a small Syberian water reservoir. *Sibirskii Ekologicheskii Zhurnal*. 2012;4:485–495. Russian.
14. Fedonenko EV, Filippova EV. Contamination level estimation in the Zaporozhian reservoir by heavy metals. *Uchenye zapiski Tavricheskogo natsional'nogo universiteta imeni V I. Vernadskogo. Seriya: Biologiya. Khimiya*. 2008;21(2):133–138. Russian.
15. Petuhova NN, Kuznetsov VA. [To Clarke values of trace elements in the soil cover of Belarus]. *Doklady Akademii nauk Belarusi*. 1992;36(5):461–465. Russian.
16. Kruchek SA, editor. *Stratigraphic charts of precambrian and phanerozoic deposits of Belarus: explanatory note*. Minsk: Belarusian Research Geological Exploration Institute; 2010. 282 p. Russian.
17. Matveev AV, Bordon VE. *Geokhimiya chetvertichnykh otlozhenii Belarusi* [Geochemistry of quaternary sediments of Belarus]. Minsk: Belaruskaja navuka; 2013. 191 p. Russian.
18. Zhuhovickaya AL, Generalova VA. *Geokhimiya ozer Belorussii* [Geochemistry of Belarussian lakes]. Minsk: Nauka i tekhnika; 1991. 202 p. Russian.
19. Petuhova NN. *Geokhimiya pochv Belorusskoi SSR* [Geochemistry of soils of the Belorussian SSR]. Minsk: Nauka i tekhnika; 1987. 231 p. Russian.
20. Lukashov KI, editor. *Geokhimicheskie provintsii pokrovnykh otlozhenii BSSR* [Geochemical provinces of superficial deposits of the BSSR]. Minsk: Nauka i tekhnika; 1969. 473 p. Russian.
21. Lukashov OV, Zhukovskaya NV, Lukashova NG, Tvoronovich-Sevruk DL, Savchenko SV. Chemical elements contents in modern bottom sediments of Belarus rivers (technochemical aspect). *Prirodopol'zovaniye*. 2015;27:109–117. Russian.
22. Farhat HI, Aly W. Effect of site on sedimentological characteristics and metal pollution in two semi-enclosed embayments of great freshwater reservoir: Lake Nasser, Egypt. *Journal of African Earth Sciences*. 2018;141:194–206. DOI: 10.1016/j.jafrearsci.2018.02.012.

Received by editorial board 11.04.2019.