DIFFERENTIABILITY PROPERTIES OF CONE-CONVEX VECTOR-VALUED FUNCTIONS

E. Bednarczuk¹, K. Leśniewski²

¹ Systems Research Institute, Polish Academy of Sciences

Newelska 6, 01-447 Warsaw, Poland

ewa.bednarczuk@ibspan.waw.pl

² Warsaw University of Technology, ul. Koszykowa 75, 00-662 Warsaw, Poland

krzlesn@gmail.com

In this talk we present some recent results on directional and Gateaux differentiability of cone-convex and cone-paraconvex vector-valued functions.

ON THE APPROXIMATION OF SYSTEMS WITH DELAY AND THEIR STABILITY

I.M. Cherevko, L.A. Piddubna, S.A. Ilika

Chernivtsi National University

Kotsyubinskogo 2, 58012 Chernivtsi, Ukraine

i.cherevko@chnu.edu.ua, laranpidd@gmail.com, ilika.svitlana17@gmail.com

Introduction. Location of the roots of the corresponding quasipolynomials plays an important role in the investigation of stability differential equations with delay. Unfortunately, we still have no effective algorithm for finding zeros of quasipolynomials.

In the present paper, we consider applications of the approximation schemes for differential-difference equations to approximate finding nonasymptotic roots of quasipolynomials, and analysis of stability for solutions of system of linear differential equations with delay.

1. Approximation schemes. Considered a linear system of differential equations with many delays

$$\frac{dx(t)}{dt} = \sum_{i=0}^k A_i x(t - \tau_i), \quad (1)$$

where $x \in \mathbb{R}^n$, A_i , $i = \overline{1, k}$, $-n \times n$ are constant matrices, $0 = \tau_0 < \tau_1 < \dots < \tau_k = \tau$.

The quasipolynomial for system (1) is of the form

$$\Phi(\lambda) = \det(\lambda E - \sum_{i=1}^k A_i e^{-\lambda \tau_i}). \quad (2)$$

In accordance with the scheme [1–3], we associate with equation (1) the system of ordinary differential equations

$$\begin{aligned}\frac{dz_0(t)}{dt} &= \sum_{i=0}^k A_i z_{l_i}(t), \quad l_i = \left[\frac{\tau_i m}{\tau} \right] \\ \frac{dz_i(t)}{dt} &= \mu [z_{i-1}(t) - z_i(t)], \quad i = \overline{1, m}, \quad \mu = \frac{m}{\tau}, \quad m \in N.\end{aligned}\tag{3}$$

For the characteristic equation of approximate system (3), we have the following equality [2]

$$\Psi_m(\lambda) = \det(\lambda E - \sum_{i=0}^k A_i (\frac{\mu}{\mu + \lambda})^{l_i}) (\mu + \lambda)^{mn}.\tag{4}$$

Moreover, the sequence of functions $H_m(\lambda) = \frac{\Psi_m(\lambda)}{(\mu + \lambda)^{mn}}$ converges as $m \rightarrow \infty$ to quasipolynomial (2). This property we can use for approximate finding nonasymptotic roots of quasipolynomial (2). Since zeros of functions $\Psi_m(\lambda)$ and $H_m(\lambda)$, we can take the zeros of characteristic polynomial (4) as approximate values on nonasymptotic zeros of quasipolynomial (2).

Theorem 1. [2] *If zero solution of equation (1) is exponentially stable (not stable) then there exists $m_0 > 0$ such that for all $m > m_0$, zero solution of system (3) is exponentially stable (not stable). If for all $m > m_0$ zero solution of approximation system (3) is exponentially stable (not stable) then zero solution of equation (1) is exponentially stable (not stable).*

2. Application. Using theorems we can obtain an effective algorithm for stability analysis of the system

$$\frac{dx}{dt} = Ax(t) + Bx(t - \tau),\tag{5}$$

where $x \in \mathbb{R}^n$, A, B — $n \times n$ are fixed matrices, $\tau > 0$.

When evaluating zeros of the characteristic equation of the approximating system of ordinary differential equations for (5) with different values of τ remaining stability of zero solution of the approximating system, we find the delay domain τ , making system (5) to be exponentially stable.

In the case where system with delay (5) is of the second order, this algorithm is easy to apply. Numerical experiments show us that system

(5) with matrices

$$A = \begin{pmatrix} -0,9 & -6,5 \\ 4,8 & -0,9 \end{pmatrix}, \quad B = \begin{pmatrix} -1,39 & -0,65 \\ 0,48 & -1,39 \end{pmatrix}$$

is asymptotically stable if $\tau \in (0, \tau_1) \cup (\tau_2, \tau_3)$, where $\tau_1 = 0,2862$, $\tau_2 = 0,7141$, $\tau_3 = 1,2142$.

References

1. *Repin Y.M.* About approximation replacement system with delay by ordinary differential equations // APM. 1965. Vol. 29, No. 2. P. 226–245.
2. *Matviy O.V., Cherevko I.M.* About approximation of system with delay and them stability // Nonlinear oscilations. 2004. Vol. 7, No. 2. P. 208–216.
3. *Ilika S.A., Matviy O.V., Pidubna L.A., Cherevko I.M.* Approximation of differential-functional equations and their application // Bukovinian Mathematical Journal. 2014. Vol. 2, No. 2–3. P. 92–96.

METHOD OF CODIFFERENTIAL DESCENT FOR GLOBAL D.C. OPTIMIZATION

M.V. Dolgopolik

Saint Petersburg State University

7/9 Universitetskaya Ebn., 199034 Saint Petersburg, Russia

maxim.dolgopolik@gmail.com

The method of codifferential descent was developed by professor V.F. Demyanov for solving a large class of nonsmooth nonconvex optimization problem in the mid-1990s [1]. Later on, several modifications of this method designed specifically for solving convex and d.c. optimization problems were proposed [2–4].

Recall that a function $f: \mathbb{R}^d \rightarrow \mathbb{R}$ is called *codifferentiable* [1] at a point $x \in \mathbb{R}^d$, if there exist compact convex sets $\underline{df}(x), \bar{df}(x) \subset \mathbb{R}^{d+1}$ such that $\max_{(a,v) \in \underline{df}(x)} a = \min_{(b,w) \in \bar{df}(x)} b = 0$, and for any $\Delta x \in \mathbb{R}^d$ one has

$$f(x+\Delta x) - f(x) = \max_{(a,v) \in \underline{df}(x)} (a + \langle v, \Delta x \rangle) + \min_{(b,w) \in \bar{df}(x)} (b + \langle w, \Delta x \rangle) + o(\Delta x),$$

where $\langle \cdot, \cdot \rangle$ is the inner product in $\mathbb{R}^d$, and $o(\alpha \Delta x)/\alpha \rightarrow 0$ as $\alpha \rightarrow +0$. The pair $Df(x) = [\underline{df}(x), \bar{df}(x)]$ is referred to as a *codifferential* of f at x . Clearly, a codifferential of f at x is not unique.

Let a function $f: \mathbb{R}^d \rightarrow \mathbb{R}$ be codifferentiable. One can utilize the method of codifferential descent (MCD) in order to minimize this function. The scheme of this method is as follows.