

## PHOTOPHYSICAL AND PHOTOCHEMICAL PROPERTIES OF HITC INDOTRICARBOCYANINE DYE MOLECULES IN SOLUTIONS

M. P. Samtsov,<sup>a\*</sup> S. A. Tikhomirov,<sup>b</sup> L. S. Lyashenka,<sup>c</sup>  
D. S. Tarasau,<sup>c</sup> O. V. Buganov,<sup>b</sup> V. A. Galievsky,<sup>b</sup>  
A. S. Stasheuski,<sup>b</sup> and E. S. Voropay<sup>c</sup>

UDC (535.37+535.34):541.14

*Results of studies on the photonics of HITC indotricarbocyanine dye molecules with  $I^-$ ,  $Br^-$ , and  $ClO_4^-$  counterions were presented. Introduction of an additional salt (tetrabutylammonium bromide) ( $10^{-2}$  M) to slightly polar solutions of the tricarbocyanine dye was shown to cause significant changes in the dye photophysical and photochemical characteristics. It was concluded based on a comparison of HITC dye photophysical properties with  $I^-$ ,  $Br^-$ , and  $ClO_4^-$  counterions in slightly polar dichlorobenzene and highly polar EtOH; an analysis of trends in HITC photobleaching in deoxygenated and air-saturated solutions; and data from femtosecond spectroscopy that the features of the photophysical and photochemical processes in slightly polar dichlorobenzene were due to superfast charge transfer in tight ion pairs followed by formation of free radicals.*

**Keywords:** indotricarbocyanine dyes, tight ion pairs, spectral properties, femtosecond transient absorption spectroscopy, photobleaching, singlet oxygen.

**Introduction.** Cationic polymethine dyes are widely used as various biological labels [1]. It seemed promising to use them as photosensitizers for cancer photochemotherapy [2, 3]. The phototoxicity mechanism of this class of compounds goes beyond a scheme in which tumor cells are damaged exclusively by the formation of singlet oxygen [3]. Indotricarbocyanine dyes were found to be localized in cancer cells in regions of low dielectric permeability and were present primarily as tight ion pairs [4, 5]. This circumstance and data that indotricarbocyanine dyes retain their photocytotoxicity under hypoxic conditions [2] suggested that photo-induced formation of free radicals as a result of charge transfer in the tight ion pairs was the starting mechanism for cancer cell damage. The occurrence of charge transfer in tight ion pairs with the formation of free radicals was confirmed by studies of transient absorption spectra of HITC dye with iodide ( $I^-$ ) and perchlorate ( $ClO_4^-$ ) counterions in the slightly polar solvent *o*-dichlorobenzene (DCB) and polar EtOH [6]. The transient spectrum of the dye in DCB was interpreted as a superposition of the absorption spectrum from the excited electronic state of the neutral molecules and the absorption spectrum of free radicals that formed practically instantaneously [6]. The high formation rate of the radicals ( $k \geq 0.6 \cdot 10^{13} \text{ s}^{-1}$ ) and the rather long lifetime of the  $S_1$ -state of the studied compound (1.8 ns) were explained by assuming that rapid phototransfer of an electron from the anion to the HITC dye cation and formation of  $An^\bullet$  and  $HITC^\bullet$  free radicals occurred only for a small part of the (tight) ion pairs.

The spectral properties of identical ion pairs can vary slightly depending on the solvent and can have different average distances between the ions [7]. Tight ion pairs with various distances between the ions are in dynamic equilibrium in an actual solution. Such pairs have similar spectral properties but can differ radically with respect to the possibility of electron transfer. It must be noted that the concept of penetrating ion pairs in which the volume of the ion pair is much less than the sum of the volumes of its component ions was introduced for polymethine dyes with specific bulky counterions in order to interpret results from phototransfer of charge in tight ion pairs [8].

Herein we present results from studies of the spectral-kinetic and photochemical properties of HITC dye under conditions where the ionic equilibrium was shifted toward an increased fraction of tight ion pairs by adding an additional salt to the solution.

\*To whom correspondence should be addressed.

<sup>a</sup>A. N. Sevchenko Institute of Applied Physical Problems, Belarusian State University, 7 Kurchatov Str., Minsk, 220108, Belarus; e-mail: samtsov@bsu.by; <sup>b</sup>B. I. Stepanov Institute of Physics, National Academy of Sciences of Belarus, Minsk; <sup>c</sup>Belarusian State University, Minsk. Translated from Zhurnal Prikladnoi Spektroskopii, Vol. 80, No. 2, pp. 177–182, March–April, 2013. Original article submitted November 29, 2012.

**Experimental.** The dye 1,3,1',3',3'-hexamethyl-2,2'-indotricarbocyanine (HITC) in three forms with different counterions ( $\text{ClO}_4^-$ ,  $\text{I}^-$ , and  $\text{Br}^-$ ) that were designated HITC( $\text{ClO}_4$ ), HITC(I), and HITC(Br) was studied. The purity of the dyes was checked by high-performance liquid chromatography (HPLC) and was also confirmed by the fact that their fluorescence spectra in EtOH were independent of excitation wavelength and their fluorescence excitation spectra were independent of recording wavelength.

The solvents EtOH and DCB were purified by standard methods [9]. An important factor in choosing the solvents was their practically identical viscosities at room temperature because the rate constants of intramolecular dissipation of electronic excitation energy in polymethine dyes depend on the medium viscosity and temperature [10, 11]. The photophysical properties of these dyes should be compared with identical solution viscosity and temperature-controlled samples.

Fluorescence and fluorescence excitation spectra and the degree of fluorescence polarization were recorded using a Fluorolog spectrofluorimeter (Spex); electronic absorption spectra, using a PV 1251A (Solar) or Specord M40 spectrophotometer. Fluorescence decay kinetics of the dyes in the nanosecond range were analyzed using a pulsed spectrofluorometer operating in the Spectroscopy Laboratory of IAPP [12]. The photodestruction quantum yield ( $\Phi$ ) of the dyes was determined by the literature method [13]. Solutions were degassed by multiple freeze–thaw cycles using a vacuum pump (down to  $10^{-3}$  Torr).

Luminescence of singlet oxygen near 1270 nm was recorded using an apparatus constructed in the Laboratory of Molecular Photonics, IP, NAS, Belarus [14]. Radiation from a semiconducting laser with  $\lambda = 667$  nm was used to excite the dyes. The experimental method was described before in detail [15]. The quantum yield of singlet-oxygen formation  $\gamma_\Delta$  was determined by a relative method. The standards were 5,10,15,20-tetrakis(4-N-methylpyridyl)porphyrin in EtOH ( $\gamma_\Delta = 0.77 \pm 0.04$ ) [16] and tetraphenylporphine (TPP) in DCB ( $\gamma_\Delta = 0.74 \pm 0.05$ ). It is noteworthy that the emission lifetime of singlet oxygen was determined during the measurements with photosensitization by the standards and experimental compounds. The singlet-oxygen lifetime in DCB was  $60 \pm 2$   $\mu\text{s}$ ; in EtOH,  $12 \pm 1$   $\mu\text{s}$ .

Transient absorption spectra were recorded using a femtosecond spectrometer [17] based on a Ti:sapphire original generator of femtosecond pulses [18] that was pumped synchronously by a pulsed Nd:YAG laser in passive synchronization mode and negative feedback. This system allowed pulses of length  $\sim 150$  fs and energies up to 1 mJ to be tuned in the range 760–820 nm at repetition rate 10 Hz. Pulses of the main frequency that were produced at the amplifier output were divided into two parts with intensity ratio 1:4. The more intense beam passed through a controlled delay line and was used as the pump pulse. Changes of the optical density of the samples were recorded using a supercontinuum generated in a 1-cm cuvette with distilled  $\text{H}_2\text{O}$  by focusing on it the smaller part of the main frequency pulse. The resulting continuum allowed a probe signal in the range 350–1000 nm to be formed taking into account the spectral sensitivity of the photomultiplier. Radiation of the continuum was divided by a semi-transparent mirror into two pulses of identical intensity (reference and signal) that were focused in the sample using an optical mirror. The reference pulse passed through the cuvette with the test compound before passage of the exciting pulse. The signal pulse passed through the excited volume with a certain delay relative to the exciting pulse. Spectra of both pulses for each laser flash were recorded by a system based on a polychromator and CCD-array with subsequent data processing.

The half-width of the cross-correlation function was  $\sim 350$  fs for all probe wavelengths. The minimal recorded change of optical density for each delay-line position with signal averaging over 100 pulses was  $\sim 0.5 \cdot 10^{-3}$ . The dye concentrations in samples for studies using the femtosecond spectrometer were  $4 \cdot 10^{-6}$ – $10^{-5}$  M. Measurements were made in 2-mm cuvettes. The optical densities of the samples were adjusted to be identical at the excitation wavelength in order to ensure similar measurement conditions. The energy of the exciting pulses was  $\sim 200$   $\mu\text{J}$  with pump beam diameter in the overlap region of the exciting and probe pulses of 6 mm. Band intensities and shapes of transient absorption spectra before the start and after the finish of a series of measurements did not differ, which indicated that the compounds were photostable.

**Results and Discussion.** The studies were carried out in EtOH and DCB for HITC( $\text{ClO}_4$ ), HITC(I), and HITC(Br), and also for HITC(Br) with tetrabutylammonium bromide (TBAB) added to the solution. This was done in order to determine if the  $\text{ClO}_4^-$  and  $\text{I}^-$  counterions were displaced from the dye molecules by  $\text{Br}^-$  upon adding the additional salt.

It was found that adding TBAB to EtOH solutions of HITC(Br) at concentrations up to  $10^{-2}$  M did not cause changes in absorption and fluorescence spectra of the dye. The half-width and position of the spectral maxima did not change (Fig. 1, Table 1). The fluorescence quantum yield and lifetime in addition to the quantum yield of singlet-oxygen generation and photodestruction also did not change. Therefore, adding TBAB did not affect the photophysical and photochemical parameters of the dye molecules in EtOH solutions where the dye was found as the free ions.