

The number of installed needles is linearly independent of the volume of the prostate gland. The same prostate volume can be irradiated using a different number of needles with DVH parameters that comply with the GEC / ESTRO guidelines [2]. Therefore, for dosimetric plans, it is possible to reduce the number of catheters while observing the requirements for target coverage and dose constraints.

For further evaluation, a more thorough analysis is required, presumably the width of the prostate gland and the installation of a fixed board relative to the organ.

#### BIBLIOGRAPHY

1. Hoskin, P. J. GEC/ESTRO recommendations on high dose rate afterloading brachytherapy for localised prostate cancer: An update / J. Peter Hoskin [et al.] // *Radiotherapy and Oncology*. – 2013. – Vol. 107. – P. 325–332.
2. Joiner, M. C. *Clinical Radiobiology* / Michael C. Joiner, Albert J. van der Kogel. Boca Raton, FL: CRC Press/Taylor & Francis Group, 2018. – P. 361.

### IMPACT OF 3-HYDROXYPYRIDINE DERIVATIVES ON CYTOSTATIC AND ANTIPROLIFERATIVE ACTIVITY OF ARABINOFURANOSYLCYTOSINE-5'-MONOPHOSPHATE

---

**M. Khanchevskii, E. Deusova**

*Belarusian State University, ISEI BSU,  
Minsk, Republic of Belarus  
maks.khanchevskiy@bk.ru*

The incidence of malignant neoplasms around the world is steadily increasing. That is why the search for drugs that prevent the development of tumors, the study of the laws of carcinogenesis is one of the main tasks of anti-tumor control. An important feature of tumor growth is the change in the level of free radical reactions, which is manifested in the increased antioxidant activity of tumor tissue, on the one hand, and the depletion of the antioxidant defense system of the tumor-bearing organism, on the other. The value of antioxidant activity is essential for the processes of cell proliferation, as antioxidants are involved in the regulation of cell reproduction. In this regard, it is attractive to search for substances or their combinations with antioxidants, the use of which will lead to a decrease in intoxication in the body of tumor carriers.

In the present work we studied the influence of modified nucleotide arabinofuranosylcytosine-5'-monophosphate in the form of the free acid (ara-CMP) and its salts with the synthetic derivatives 3-hydroxypyridine emoxipin (ara-CMP+Em) on the viability of mononuclear cells in the peripheral blood, the number and lymphocyte proliferation in mitogen-induced stimulation of the cells. Under ara-CMP lymphocytes not only stopped cell division and increased cell death, but as well as fractions of secreting pro-inflammatory cytokines cells were increased. It is known that increasing of pro-inflammatory cytokines level is a systemic reaction to the increased ROS levels due to destruction of cells. Note that the presence of emoxipin (substances that have antioxidant properties) almost completely neutralized the observed effect.

#### BIBLIOGRAPHY

1. Tsesmetzis, N. *Cancers* / N. Tsesmetzis, C. B. J. Paulin, S. G. Rudd, N. Herold. – 2018. – № 10. – C. 240–248.
2. Block, K. I. *Cancer* / A. C. Koch, M. N. Mead, P. K. Tothy, R. A. Newman, C. Gyllenhaal. *Int. J.* – 2008. – № 15. – C. 1227–1239.
3. Pan, H. *Pharmacol.* / H. Pan, P. Mukhopadhyay, M. Rajesh, V. Patel, B. Mukhopadhyay, B. Gao, G. Haskó, P. J. Pacher. – *Exp. Ther.* – 2008. – № 12. – C. 24–28.

#### Acknowledgement

This work was supported by the BRFFR (grant X18MB-019).