

Management of temporal parameters of super-powerful laser pulses

S. Mironov, V. Lozhkarev, E. Khazanov

Nonlinear dynamic and optics Department, Institute of Applied Physics, Nizhny Novgorod, Russia, e-mail: Mironov_s@appl.sci-nnov.ru

The report is devoted to the management of pulse duration and temporal intensity contrast ratio enhancement of laser pulses with petawatt power. The theoretical and experimental results of second harmonic generation process under strong influence of cubic nonlinearity will be discussed. The outcomes of implementation of small-scale self-focusing suppression method at intense laser beams ($1\div 4\text{TW}/\text{cm}^2$) will be considered.

WKB approach applied to 1D time-dependent nonlinear Hamiltonian oscillators

M. Robnik

CAMTP - Center for Applied Mathematics and Theoretical Physics, University of Maribor, Maribor, Slovenia, European Union, e-mail: Robnik@uni-mb.si

We present a simple WKB-like approach to obtain approximate analytic solutions to a certain class of time-dependent nonlinear 1D Hamiltonian oscillators. The case of homogeneous power law potentials is solved explicitly in closed form in the leading order. The accuracy of the approximation is surprisingly good and we illustrate it in case of the quartic oscillator. The method is expected to be useful in treating the statistical properties of the dynamical quantities (like energy and adiabatic invariant) in nonlinear oscillators, and in many other applications in nonlinear oscillators.

References:

Papamikos G and Robnik M 2012 J.Phys.A: Math.Theor. 45 0152206

Robnik M and Romanovski V G 2000 J.Phys.A: Math.Theor. 33 5093

Robnik M V and Romanovski V G 2008 "Let's Face Chaos through Nonlinear Dynamics",

Proceedings of the 7th International summer school/conference, Maribor-Slovenia, 2008, in AIP Conf. Proc. 1076, Eds. M.Robnik and V.G. Romanovski.

Chaotic dynamics of generation of radiation by relativistic electron beams in photonic crystals

S. Sytova

Research Institute for Nuclear Problems, Belarusian State University, Minsk, Belarus, e-mail: sytova@inp.bsu.by

We consider generation of radiation by relativistic electron beams in photonic (electromagnetic) crystals in conditions of Volume Free Electron Lasers (VFEL).

Such crystals include natural crystals and artificial three-dimensional periodical structures. In VFEL while relativistic electron beam passing through such photonic crystals in conditions of dynamical diffraction quasi-Cherenkov parametric radiation appears and obeys to special law [Baryshevsky V.G., Feranchuk I. D. Phys. Lett. A. 102 (1984), 141]. This law is valid for all wavelength ranges and substantially differs from its analogues in other electronic devices such as traveling wave tubes, backward-wave tubes, free electron lasers and other. Different sides of chaotic dynamics of such generation are investigated. The complicated root to chaos in VFEL is demonstrated starting with the generation threshold.

Laser pulse control of DCI^+ photofragmentation

M.V. Korolkov

IF NANB, Minsk, Belarus, e-mail: korolkovmv@yahoo.com

The prospects of the control of deuterium chloride ion (DCI^+) photofragmentation with the laser field generated by two time-shifted interfering femtosecond laser pulses is investigated theoretically. The analysis performed by means of computer simulation within the Schroedinger wave function formalism. The calculations provide evidence that the ratio of product ion yields Cl^+ versus D^+ can be manipulated by appropriate choice of laser pulse parameters, in particular central laser wavelength, pulse duration, intensity and chirp. The analysis of time dependent populations reveals competition between intra- and inter - electronic state excitation, enabling the understanding of quantum control at the molecular level.

On some peculiarities in transition from de Sitter's to Minkowski model: iteration procedure in the radial equation for a scalar particle

V.M. Red'kov, G.G. Krylov, E.M. Ovsiyuk

Institute of Physics, NAS, Belarusian State University, Mozyr State Pedagogical University, e-mail: redkov@dragon.bas-net.by

Though the problem of Hawking radiation in de Sitter space-time, in particular details of penetration of a quantum mechanical particle through the de Sitter horizon, has been examined intensively there is still some vagueness in this subject. The present paper aims to clarify the situation. A known algorithm for calculation of the reflection coefficient R_{ej} on the background of the de Sitter space-time model is analyzed. It is shown that the determination of R_{ej} requires an additional constrain on quantum numbers $\varepsilon R/\hbar c \gg j$, where R is a curvature radius. When taking into account this condition, the value of R_{ej} turns out to be precisely zero. It is shown that the basic instructive definition for the calculation