

Corrections for a constant radial magnetic field in the muon $g-2$ and electric-dipole-moment experiments in storage rings

Alexander J. Silenko^{1,2,a}

¹ Research Institute for Nuclear Problems, Belarusian State University, Minsk 220030, Belarus

² Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna 141980, Russia

Received: 30 May 2017 / Accepted: 18 September 2017 / Published online: 22 October 2017

© The Author(s) 2017. This article is an open access publication

Abstract We calculate the corrections for constant radial magnetic field in muon $g - 2$ and electric-dipole-moment experiments in storage rings. While the correction is negligible for the current generation of $g - 2$ experiments, it affects the upcoming muon electric-dipole-moment experiment at Fermilab.

1 Introduction

Muon $g - 2$ and electric-dipole-moment (EDM) experiments in storage rings are of great importance for contemporary physics. The well-known discrepancy between theoretical and experimental values of the anomalous magnetic moment (AMM) of muon [1, 2] may be a window to new physics. New physics can result in a higher value of the muon EDM than that predicted by the Standard Model.

In such precise experiments it is important to fully account for all significant corrections affecting the measurement. This is the reason for conducting the present research which determines corrections for a constant radial magnetic field in high-precision $g - 2$ and EDM experiments with polarized muon beams in storage rings. Increasing of the accuracy of the planned experiments as compared to the accomplished ones makes this research necessary.

We use the cylindrical coordinate system and the system of units with $\hbar = 1$, $c = 1$. We include \hbar and c into some equations when this inclusion clarifies the problem.

2 General equations of spin motion in the cylindrical and the Frenet–Serret coordinate systems

In this section, we briefly report the results of a comparison of the cylindrical and the Frenet–Serret coordinate sys-

tems obtained in Ref. [3]. While these coordinate systems are equivalent, the related equations of the spin motion differ. The axes of the Frenet–Serret coordinate system are motional and are defined relative to the particle velocity direction. In this system, the equation of spin motion defines a motion of the spin (pseudo)vector with respect to the momentum vector, i.e., the change in a relative orientation of these vectors. The Frenet–Serret coordinate system is commonly used in accelerator theory. For a particle with electric and magnetic dipole moments, the classical equation of the spin motion is given by (see Refs. [4, 5] and the references therein)

$$\Omega^{(FS)} = -\frac{e}{m} \left[a\mathbf{B} - \frac{a\gamma}{\gamma+1} \boldsymbol{\beta}(\boldsymbol{\beta} \cdot \mathbf{B}) + \left(\frac{1}{\gamma^2-1} - a \right) (\boldsymbol{\beta} \times \mathbf{E}) + \frac{\eta}{2} \left(\mathbf{E} - \frac{\gamma}{\gamma+1} \boldsymbol{\beta}(\boldsymbol{\beta} \cdot \mathbf{E}) + \boldsymbol{\beta} \times \mathbf{B} \right) \right], \quad (1)$$

where $a = (g - 2)/2$, $\boldsymbol{\beta} = \mathbf{v}/c$, $g = 2mc\mu/(e\hbar s)$, $\eta = 2mcd/(e\hbar s)$. Here s is the spin quantum number and the η factor for the EDM d is similar to the g one for the magnetic dipole moment.

When the cylindrical coordinate system is used, the spin motion equation is defined allowing for motion of the horizontal axes of the cylindrical coordinate system which follows the azimuthal motion of a given particle. The vertical axes of the cylindrical and the Cartesian coordinate systems always coincide. The angular velocities of the spin motion in these coordinate systems differ on $\dot{\phi}\mathbf{e}_z$, where $\dot{\phi}$ is the angular velocity of a change of the particle azimuth in the storage ring [6]. As a result, the spin motion equation in the cylindrical coordinate system takes the form [3, 6]

$$\Omega^{(cyl)} = -\frac{e}{m} \left\{ a\mathbf{B} - \frac{a\gamma}{\gamma+1} \boldsymbol{\beta}(\boldsymbol{\beta} \cdot \mathbf{B}) + \left(\frac{1}{\gamma^2-1} - a \right) (\boldsymbol{\beta} \times \mathbf{E}) + \frac{1}{\gamma} \left[\mathbf{B}_{\parallel} - \frac{1}{\beta^2} (\boldsymbol{\beta} \times \mathbf{E})_{\parallel} \right] + \frac{\eta}{2} \left(\mathbf{E} - \frac{\gamma}{\gamma+1} \boldsymbol{\beta}(\boldsymbol{\beta} \cdot \mathbf{E}) + \boldsymbol{\beta} \times \mathbf{B} \right) \right\}. \quad (2)$$

^a e-mail: alsilenko@mail.ru

The sign \parallel indicates the horizontal projection for any vector. This equation describes the spin motion relative to immobile detectors placed around the storage ring. Equation (2), contrary to Eq. (1), does not depend on the vertical motion of the beam. The compactness of Eq. (1) relies on the fact that the Frenet–Serret coordinate axes move relative to the immobile detectors. As a result, Eq. (1) needs to be used with care. In particular, this equation may create the illusion that, on condition that $\beta \cdot \mathbf{B} = 0$, the impact of vertical and radial fields on the spin is identical. For leptons (electron, muon) the difference between $\Omega^{(FS)}$ and $\Omega^{(cyl)}$ can be rather significant. To determine an *observable* effect, the motion of the tangential and normal axes of the Frenet–Serret coordinate system should be added to the spin motion in this coordinate system. Once this factor is taken into account, the cylindrical and the Frenet–Serret coordinate systems give an equivalent description of the spin motion [3].

The additional term in Eq. (2) which is proportional to $1/\gamma$ can increase the importance of the horizontal components of \mathbf{B} and $\beta \times \mathbf{E}$ as compared to Eq. (1). In particular, the spin quantization axis (also called the stable spin axis) in a purely magnetic storage ring does not coincide with the direction of the total magnetic field. In relation to the longitudinal magnetic field, this property has been considered in Ref. [7].

It should be added that the quantum-mechanical description of the spin motion for spin-1/2 [8] and spin-1 [9] particles fully agrees with the classical description.

3 Manifestations of the constant radial magnetic field in muon $g - 2$ and electric-dipole-moment experiments

The correction for the constant radial magnetic field is nonzero in both the $g - 2$ and the electric-dipole-moment experiments with polarized muons in storage rings. However, the importance of the systematic effects caused by this field significantly differs in the two cases. For a calculation of this correction, it is convenient to use Eq. (2). Since the *average* Lorentz force acting on a particle is zero, the average values of the vertical electric field and the radial magnetic one satisfy the condition

$$E_z = -(\beta \times \mathbf{B})_z. \quad (3)$$

When electric focusing is used, the angular velocity of the spin precession is given by

$$\Omega^{(cyl)} = -\frac{e}{m} \left\{ a B_z \mathbf{e}_z + \frac{g}{2\gamma^2} B_\rho \mathbf{e}_\rho + \frac{\eta}{2} (\beta \times \mathbf{B})_\parallel \right\}. \quad (4)$$

It is well known that the fields satisfying Eq. (3) turn the spin. The vertical electric field conditions are associated with one of the main systematic errors in EDM experiments in

storage rings carried out by the frozen spin method [10]. The condition (3) is satisfied for Wien filters with static [11] and oscillating [12–14] fields, which are frequently used as spin rotators. In particular, the expression for the radial component of Ω (halved owing to the field oscillations) is presented in Ref. [12].

In the muon $g - 2$ and EDM experiments described in Refs. [1, 15], $\gamma^2 = 1 + a^{-1}$ and $g/(2\gamma^2) = a$. As a result, Eq. (4) takes the form

$$\Omega^{(cyl)} = -\frac{e}{m} \left\{ a(B_z \mathbf{e}_z + B_\rho \mathbf{e}_\rho) + \frac{\eta}{2} (\beta \times \mathbf{B})_\parallel \right\}. \quad (5)$$

The angular velocity of the spin rotation caused by the magnetic moment is collinear to the vector of the total magnetic field \mathbf{B} . However, the vertical and radial components of \mathbf{B} enter Eq. (4) with different factors and the spin motion defined by Eq. (5) is influenced not only by the radial magnetic field but also by the vertical electric one.

Equation (5) can be presented as follows:

$$\begin{aligned} \Omega^{(cyl)} &= \omega_0 \mathbf{e}_z + O \mathbf{e}_\rho, \quad \omega_0 = -\frac{e}{m} a B_z, \\ O &= \omega_0 \left(\frac{B_\rho}{B_z} + \frac{\eta \beta}{2a} \right), \end{aligned} \quad (6)$$

where the quantities ω_0 , O , and β ($\beta = \beta \mathbf{e}_\phi$) can be positive and negative.

Evidently, the constant radial magnetic field conditions the following correction to the $g - 2$ frequency:

$$\Delta\omega = -\omega_0 \frac{B_\rho^2}{2B_z^2}. \quad (7)$$

In the $g - 2$ experiments [1, 2], this correction can be neglected. Measurements of the constant radial magnetic field in the BNL muon $g - 2$ experiment have shown that $|B_\rho| \approx 2.9 \times 10^{-5} \text{ T}$. Since $B_z = 1.45 \text{ T}$, $|B_\rho/B_z| \approx 2 \times 10^{-5}$ [16]. In this case, the correction to the $g - 2$ frequency is of the order of 10^{-10} and can be neglected.

However, the constant radial magnetic field can be more important for the muon EDM experiments. The search for the muon EDM has been performed in the framework of the past BNL experiment [16]. The initial beam polarization was longitudinal and the vertical spin component was detected. This experiment has established the upper bound on the muon EDM, $|d_\mu| < 1.8 \times 10^{-19} \text{ e-cm}$ [15].

In the considered case, squares of small parameters can be neglected. The spin evolution is defined by the general equation (15) in Ref. [17] and (11) in Ref. [18], which takes the form

$$\begin{aligned} P_\rho(t) &= \cos(\omega_0 t + \psi), \quad P_\phi(t) = \sin(\omega_0 t + \psi), \\ P_z(t) &= -\frac{O}{\omega_0} \cos(\omega_0 t + \psi), \end{aligned} \quad (8)$$

where ψ is equal to $\pi/2$ and $-\pi/2$ for counterclockwise and clockwise muon beams, respectively, and \mathbf{P} is the polarization vector. In this case, a nonzero vertical component of the polarization vector is conditioned by the terms *linear* in B_ρ/B_z and η . The related correction to the $g-2$ frequency is *quadratic or bilinear* in B_ρ/B_z and η .

Equations (6) and (8) demonstrate that the constant radial magnetic field can mimic an EDM. The constant radial magnetic field measured in Ref. [16] corresponds to the muon EDM of 2×10^{-21} e·cm. This value is small as compared with the current bound on the muon EDM established in Ref. [15]. However, the sensitivity to the muon EDM will be increased by two orders of magnitude in the future experiment at Fermilab [19]. In this case, the correction for a constant radial magnetic field needs to be considered. We can add that a muon EDM experiment based on the frozen spin method [10] is sensitive to the muon EDM of the order of 10^{-24} e·cm.

4 Summary

In the present paper, we have calculated the correction for the constant radial magnetic field in the $g-2$ and electric-dipole-moment experiments with muons in storage rings. While the correction can be neglected in the $g-2$ experiments [1,2], it is important for the forthcoming muon EDM experiment at Fermilab.

The planned $g-2$ /EDM experiment at J-PARC will be performed in a purely magnetic ring [20,21]. In this case, the average constant radial magnetic field is equal to zero and the correction for this field vanishes.

Acknowledgements The author is grateful to N. Nikolaev for useful comments and discussions. The author acknowledges the support by the Belarusian Republican Foundation for Fundamental Research (Grant No. $\Phi 16D-004$) and by the Heisenberg-Landau program of the German Ministry for Science and Technology (Bundesministerium für Bildung und Forschung).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

References

1. G.W. Bennett et al., (Muon $g-2$ Collaboration), Final report of the E821 muon anomalous magnetic moment measurement at BNL. Phys. Rev. D **73**, 072003 (2006)
2. A. Chapelain, The Muon $g-2$ experiment at Fermilab. EPJ Web. Conf. **137**, 08001 (2017)
3. A.J. Silenko, Comparison of spin dynamics in the cylindrical and Frenet-Serret coordinate systems. Phys. Part Nucl. Lett. **12**, 8 (2015)
4. T. Fukuyama, A.J. Silenko, Derivation of generalized Thomas-Bargmann-Michel-Telegdi equation for a particle with electric dipole moment. Int. J. Mod. Phys. A **28**, 1350147 (2013)
5. A.J. Silenko, Spin precession of a particle with an electric dipole moment: contributions from classical electrodynamics and from the Thomas effect. Phys. Scr. **90**, 065303 (2015)
6. A.J. Silenko, Equation of spin motion in storage rings in the cylindrical coordinate system. Phys. Rev. ST Accel. Beams **9**, 034003 (2006)
7. A.J. Silenko, Polarization of spin-1/2 particles in an axisymmetric magnetic field. JETP **87**, 629 (1998)
8. A.J. Silenko, Quantum-mechanical description of the electromagnetic interaction of relativistic particles with electric and magnetic dipole moments. Russ. Phys. J. **48**, 748 (2005)
9. A.J. Silenko, Quantum-mechanical description of spin-1 particles with electric dipole moments. Phys. Rev. D **87**, 073015 (2013)
10. F.J.M. Farley, K. Jungmann, J.P. Miller, W.M. Morse, Y.F. Orlov, B.L. Roberts, Y.K. Semertzidis, A. Silenko, E.J. Stephenson, A new method of measuring electric dipole moments in storage rings. Phys. Rev. Lett. **93**, 052001 (2004)
11. V. Tioukine, K. Aulenbacher, Operation of the MAMI accelerator with a Wien filter based spin rotation system. Nucl. Instrum. Methods Phys. Res. A **568**, 537 (2006)
12. W.M. Morse, Y.F. Orlov, Y.K. Semertzidis, rf Wien filter in an electric dipole moment storage ring: the partially frozen spin effect. Phys. Rev. ST Accel. Beams **16**, 114001 (2013)
13. J. Slim, R. Gebel, D. Heberling, F. Hinder, D. Hölscher, A. Lehrach, B. Lorentz, S. Mey, A. Nass, F. Rathmann, L. Reifferscheidt, H. Soltner, H. Straatmann, F. Trinkel, J. Wolters, Electromagnetic simulation and design of a novel waveguide RF Wien filter for electric dipole moment measurements of protons and deuterons. Nucl. Instrum. Methods A **828**, 116 (2016)
14. A. Saleev et al. (JEDI Collaboration), Spin tune mapping as a novel tool to probe the spin dynamics in storage rings. Phys. Rev. ST Accel. Beams **20**, 072801 (2017)
15. G.W. Bennett et al. (Muon $g-2$ Collaboration), Improved limit on the Muon electric dipole moment. Phys. Rev. D **80**, 052008 (2009)
16. S.I. Redin, N.M. Ryskulov, G.V. Fedotov, B.I. Khazin, G.M. Bunce, G.T. Danby, J.W. Jackson, W.M. Morse, R. Prigl, Y.K. Semertzidis, E. Efstathiadis, B.L. Roberts, A. Grossmann, K. Jungmann, G. zu Putlitz, P. von Walter, S.K. Dhawan, F.J.M. Farley, M. Grosse-Perdekamp, V.W. Hughes, D. Kawall, Radial magnetic field measurements with a Hall probe device in the muon ($g-2$) storage ring magnet at BNL. Nucl. Instrum. Methods Phys. Res. A **473**, 260 (2001)
17. A.J. Silenko, General classical and quantum-mechanical description of magnetic resonance: an application to electric-dipole-moment experiments. Eur. Phys. J. C **77**, 341 (2017)
18. A.J. Silenko, General description of spin motion in storage rings in the presence of oscillating horizontal fields. Europhys. Lett. **118**, 61003 (2017)
19. R. Chislett on behalf of the Muon $g-2$ Collaboration, The muon EDM in the $g-2$ experiment at Fermilab. EPJ Web Conf. **118**, 01005 (2016)
20. T. Mibe, the J-PARC collaboration, Measurement of muon $g-2$ and EDM with an ultra-cold muon beam at J-PARC. Nucl. Phys. B Proc. Suppl. **218**, 242 (2011)
21. M. Otani (for the E34 collaboration), Status of the Muon $g-2$ /EDM Experiment at J-PARC (E34). JPS Conf. Proc. **8**, 025008 (2015)