

First Record of the Black Locust Gall Midge *Obolodiplosis robiniae* (Haldeman, 1847) (Diptera: Cecidomyiidae) in Türkiye

Aleh V. SINCHUK^{1, 2*}

Nadzeya V. SINCHUK³

¹Scientific and Practical Center of the National Academy of Sciences of Belarus for Bioresources, Laboratory of Terrestrial Invertebrate Animals, Minsk 220072, BELARUS

²Belarusian State University, Faculty of Geography and Geoinformatics, UNESCO Chair in Science Education with Emphasis on Natural Sciences, Minsk 220030 BELARUS

³Vitebsk region, Polotsk district, Gorodishche vill., 21A 211658 BELARUS

e-mails: ^{1,2}aleh.sinchuk@gmail.com, ³n.v.sinchuk@gmail.com

ORCID IDs: ¹0000-0003-2845-1180, ³0000-0002-4116-7283

*Corresponding author

ABSTRACT

In 2019 galls of black locust gall midge *Obolodiplosis robiniae* (Haldeman, 1847) (Diptera: Cecidomyiidae) were discovered in Türkiye on the black locust tree, *Robinia pseudoacacia* L. (Fabaceae) in Üçkardeş village (Kemalpaşa District, Artvin Province). The black locust gall midge is a new invasive species in Türkiye. *Obolodiplosis robiniae* was spread from Georgia.

Keywords: invasive species, *Robinia pseudoacacia*, leaf damage, phytophagous insects, species distribution, new data.

Sinchuk, A. V. & Sinchuk, N. V. (2025). First record of the black locust gall midge *Obolodiplosis robiniae* (Haldeman, 1847) (Diptera: Cecidomyiidae) in Türkiye. *Journal of the Entomological Research Society*, 27(2), 307-313.

Received: 13 February, 2025

Accepted: July 08, 2025

INTRODUCTION

The black locust (*Robinia pseudoacacia* L.) is an alien species in Europe. It was introduced in the 17th century from North America to Europe in order to consolidate soils and to reforest devastated areas, or as an ornamental tree (Sitzia, Cierjacks, De Rigo & Caudullo, 2016). Various phytophagous insect species are associated with the black locust, such as the leaf miners *Parectopa robinella* (Clemens, 1863) (Vidano, 1970), *Macrosaccus robinella* (Clemens, 1859) (Whitebread, 1989), locust sawfly *Nematus tibialis* Newman, 1837 (Maslyakov & Izhevsky, 2011), black locust gall midge *Obolodiplosis robiniae* (Haldeman, 1847) (Duso & Skuhravá, 2003) and its specialized parasitoid *Platygaster robiniae* Buhl & Duso, 2007 (Buhl & Duso, 2008) has been accidentally introduced to Europe.

The gall midge *Obolodiplosis robiniae*, like its host a native to North America, was first reported in Europe in 2003 in the Veneto region in Italy (Duso & Skuhravá, 2003). Then it has been spread to Czech Republic (Skuhravá & Skuhřavý, 2005; Šefrova & Lantuvka, 2005), Slovenia (Duso, Fontana, & Tirello, 2005), Croatia (Skuhravá, Skuhřavý, & Csóka, 2007), Germany (Wehrmaker, 2007), Hungary (Csóka, 2006), Slovakia (Tóth, Váňová, & Lukáš, 2009), Ukraine (Berest, 2006), Montenegro (Glavendekić, Mihajlović, Jakovljević, & Marjanović, 2008), Netherlands (Roskam, van der Aa, Bijkerk, Ellis, & Moraal, 2008), Switzerland (Wermelinger & Skuhravá, 2007), Austria (Tomiczek, Cech, Fürst, Hoyer-Tomiczek, Krehan, Perny, & Steyrer, 2009), France (Laguerre & Dauphin, 2007), Poland (Skrzypczyńska, 2007), Serbia (Mihajlović, Glavendekić, Jakovljević, & Marjanović, 2008), United Kingdom (Skuhravá & Skuhřavý, 2005), Luxemburg (Schneider & Walisch, 2009), Sweden (Molnár, Boddum, Szöcs, & Hillbur, 2009), Denmark (Jørgensen, 2009), Bulgaria (Tomov et al., 2009), Croatia (Pernek & Matošević, 2009), Romania (Bálint, Neacșu, Balog, Fail, & Véték, 2010), Belarus (Petrov, 2010), Moldova (Mocreac & Timuș, 2014), Russia (Gninenko, Kostukov, & Kosheleva, 2011; Sinchuk, Roginsky, & Buga, 2019), Lithuania (Stalažs, 2014), Albania, Macedonia, Bosnia and Herzegovina (Skuhravá, Martinez, & Roques, 2010), Portugal (Bella, 2013), Spain (Sánchez & Umaran, 2013), Greece (Bella, 2014), Latvia (Rupais, Stalazs, & Strelcuns, 2014). The black locust gall midge was found in Georgia in 2013 (Skuhravá, Skuhřavý, & Buhr, 2013). The latest European report is from Armenia, where the species was detected in 2019 (Gubin, 2021). The attack occurs in the species *R. pseudoacacia* L., *R. viscosa* L. and *R. hispida* L., mainly in urban areas, where it mainly affects their aesthetic value. The larvae develop characteristic leaf margin roll galls on the infested leaves. There are up to 6 galls on a leaflet, and on average there are up to 5-6 larvae per gall. In Europe it develops two, three and, in optimal conditions, even four generations a year (Ilea, Hulujan, Florian, Florian, & Oltean, 2023). The current potential distribution area of *O. robiniae* is within the range of 21,57 and 65,65 N in the Eurasian continent. The primary factor controlling the distribution of *O. robiniae* is temperature. The highly and moderately suitable areas are mainly distributed in the semi-humid and semi-arid regions, which also happen to be the locations where the host black locust (*Robinia pseudoacacia* L.) grows at its fastest rate (Zhao, Gao, Du, & Shi, 2023). The forecast

of the potential distribution area of *O. robiniae* revealed that the species would benefit from global warming. The assessment of climate conditions has revealed that the Black Sea coast, as well as the northwestern and northeastern regions of Türkiye, are highly suitable for the settlement of the black locust gall midge. In contrast, the southeastern part of Türkiye is deemed moderately suitable for this invasive insect (Zhao, Gao, Du, & Shi, 2023).

Currently, 121 species of gall midges (Cecidomyiidae) have been recorded in Turkey. Representatives of the genus *Obolodiplosis* have not been found to date (Skuhravá, Bayram, Çam, Tezcan, & Can, 2005; Cilbircioğlu & Ünal, 2008; Tataroğlu, Katılmış, & Skuhravá, 2024).

MATERIALS AND METHODS

This article is based on the material of damaged leaves, which were found on *R. pseudoacacia*. Damaged rolled leaves were collected in the north-eastern part of Türkiye in coastal Üçkardeş village (Kemalpaşa District, Artvin Province) (Karadeniz Sahil Yolu, 41.490127N, 41.530488E, 08.10.2019, leg. A.V. Sinchuk). The number of damaged leaves in relation to not damaged in the lower tree crown was estimated and the number of damaged leaves in a complex leaf was established.

The damaged leaf blades were collected in Zip-Lock plastic bags in order to eliminate moisture loss and subsequent drying of the material. The damaged leaf blades were herbarized. Identification of the pests was carried out by using specialized keys by damaged leaf blades (Ellis, 2001–2025).

RESULTS AND DISCUSSION

Examination of characteristic leaf damage led to the identification of the dipteran species – *Obolodiplosis robiniae*. The occupancy of leaf blades of *R. pseudoacacia* by black locust gall midge in the lower tree crown was up to 5 %. Complex leaves were marked with 1–2 damaged leaves. The black locust gall midge *O. robiniae* has spread quite fast from the place of its first record in Europe.

A low density of *O. robiniae* was noticed. The black locust gall midge is a new invasive species in Türkiye. The dispersal in Türkiye has one main entry from Georgia. However, the records of the border Georgia show its wide distribution (Skuhravá et al., 2013). The black locust gall midge was discovered in the Botanical Garden in Batumi. The pest density was not high – the occupancy was 2 %. Complex leaves were marked with one damaged leaf. Türkiye is a global potential distribution area for black locust on the same level as Georgia (Li, Xu, Guo, & Du, 2014).

Georgia is a spatial abundance of black locust specimens. It can be argued that along with the spread of the plant, the spread of the pest occurs due to the high traffic stream and natural expansion is possible too. *O. robiniae* could have entered Turkey through natural range expansion from Batumi (Georgia).

R. pseudoacacia is a species introduced to Europe from North America. The main production areas of black locust plants for export are Istanbul, Bursa, and Sakarya, with additional production zones in Adana, Antalya, and Izmir (EFSA PLH Panel, 2021). According to GBIF, *R. pseudoacacia* is recorded throughout Turkey along the Black Sea and Mediterranean coastlines (GBIF, 2025). These areas represent potential habitats for the establishment of *O. robiniae*.

According to models predicting the future distribution of the species in Eurasia for the years 2050 and 2070, the black locust gall midge is expected to expand its range across Eastern and parts of Central Türkiye (Zhao et al., 2023). Further research is essential to accurately assess the current distribution status of *O. robiniae*.

CONCLUSIONS

The black locust gall midge (*Obolodiplosis robiniae* (Haldeman, 1847) were discovered in Türkiye (Üçkardeş village, Kemalpaşa District, Artvin Province). The black locust gall midge larvae feed on the leaf blades of *Robinia pseudoacacia*. Low pest density was observed ($\leq 5\%$ leaf damage in lower crowns), consistent with early-stage colonization. Proximity to Batumi, Georgia – where *O. robiniae* is established – supports natural range expansion as the primary introduction route. Human-mediated dispersal (e.g., trade, transport) remains plausible but unverified. Coastal regions (Black Sea, Mediterranean) and commercial *R. pseudoacacia* production zones (Istanbul, Bursa, Sakarya) are highly suitable habitats for establishment. Climate models predict range expansion into eastern/central Türkiye by 2050–2070 under warming scenarios. The gall midge fauna of Turkey comprises 122 species (Cecidomyiidae).

REFERENCES

- Bălint, J., Neacșu, P., Balog, A., Fail, J., & Véték, G. (2010). First record of the black locust gall midge, *Obolodiplosis robiniae* (Haldeman) (Diptera: Cecidomyiidae), in Romania. *North-Western Journal of Zoology*, 6(2), 319–322.
- Bella, S. (2013). New alien insect pests to Portugal on urban ornamental plants and additional data on recently introduced species. *Annales de la Société Entomologique de France*, 49(4), 374–382. <https://doi.org/10.1080/00379271.2013.856210>.
- Bella, S. (2014). Invasive insect pests and their associated parasitoids on ornamental urban plants on Corfu island – *Phytoliriomyza jacarandae* Steyskal and Spencer 1978 (Diptera, Agromyzidae) a new record in Greece. *Hellenic Plant Protection Journal*, 7(2), 53–59.
- Berest, Z. L. (2006). Obnaruzheniye gallitsy *Obolodiplosis robiniae* (Diptera, Cecidomyiidae) v Ukraine [The discovery of the gall-midges *Obolodiplosis robiniae* (Diptera, Cecidomyiidae) in Ukraine]. *Vestnik Zoologii*, 40(6), 534 (in Russian).
- Buhl, P. N. & Duso, C. (2008). *Platygastr robiniae* n. sp. (Hymenoptera: Platygastridae) parasitoid of *Obolodiplosis robiniae* (Diptera: Cecidomyiidae) in Europe. *Annals of the Entomological Society of America*, 101(2), 297–300.
- Cilbiricioğlu, C. & Ünal, S. (2008). Gall Midges (Diptera: Cecidomyiidae) in forest trees of Turkey. *Journal of Agricultural and Urban Entomology*, 25(1), 13–23. <https://doi.org/10.3954/1523-5475-25.1.13>.

First Record of the Black Locust Gall Midge Obolodiplosis robiniae

- Csóka, G. (2006). Az akác-gubacsszúnyog [*Obolodiplosis robiniae* (Haldeman 1847)] megjelenése Magyarországon [Appearance of the black locust gall midge [*Obolodiplosis robiniae* (Haldeman 1847)] in Hungary]. *Növényvédelem*, 42(12), 663-665 (in Hungarian).
- Duso, C. & Skuhrová, M. (2003). First record of *Obolodiplosis robiniae* (Haldeman) (Diptera Cecidomyiidae) galling leaves of *Robinia pseudoacacia* L. (Fabaceae) in Italy and Europe. *Frustula Entomologica*, 25(38), 117-122.
- Duso, C., Fontana, P., & Tirello, P. (2005). Diffusione in Italia e in Europa di *Obolodiplosis robiniae* (Haldeman), dittero cecidomide nearctico dannoso a *Robinia pseudoacacia* [Spread of the gall midge *Obolodiplosis robiniae* (Haldeman) injurious to black locust in Italy and Europe]. *Informatore Fitopatologico*, 55(5), 30-33 (in Italian).
- Ellis, W.N. (2001–2025). *Leafminers and plant galls of Europe*. Retrieved from <https://bladmineerders.nl> (last modified 9 January 2020; accessed 3 January 2025).
- EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Dehnen-Schmutz, K., Di Serio, F., Gonthier, P., Jacques, M.-A., Jaques Miret, J.A., Justesen, A.F., MacLeod, A., Magnusson, C.S., Milonas, P., Navas-Cortes, J.A., Parnell, S., Reignault, P.L., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L., Chatzivassiliou, E., Debode, J., Manceau, C., Gardi, C., Mosbach-Schulz, O. & Potting, R. (2021). Scientific Opinion on the commodity risk assessment of *Robinia pseudoacacia* plants from Turkey. *EFSA Journal*, 19(5), 6568, 54 p. <https://doi.org/10.2903/j.efsa.2021.6568>.
- GBIF.org (2025). *GBIF Occurrence*. Retrieved from <https://doi.org/10.15468/dl.j93tz6> (accessed 30 June 2025).
- Glavendekić, M., Mihajlović, Lj., Jakovljević, I., & Marjanović, S. (2008). *Obolodiplosis robiniae* (Haldeman) (Diptera: Cecidomyiidae) – a new invasive insect pest in Serbia. *Glasnik Sumarskog fakulteta*, 97, 215-226.
- Gninenko, Y.I., Kostukov, V.V., & Kosheleva, O.V. (2011). Novye invazivnye nasekomye v lesah i ozelenitel'nyh posadkah Krasnodarskogo kraya [New invasive insects in the forests and greenery of the Krasnodar krai]. *Zashita i Karantin Rastenii*, (4), 49-50 (in Russian).
- Gubin, A.I. (2021). Four invasive alien phytophagous insects new to Armenia. *Phytoparasitica*, 49, 163-166. <https://doi.org/10.1007/s12600-020-00853-0>.
- Ilea, I., Hulujan, I.-B., Florian, T., Florian, V., & Oltean, I. (2023). The gall midge *Obolodiplosis robiniae* Haldemann (Diptera Cecidomyiidae) new invasive alien species in Europa – Review. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 51(2), 13096. <https://doi.org/10.15835/nbha51213096>.
- Jørgensen, J. (2009). *Obolodiplosis robiniae* (Haldeman, 1847) (Cecidomyiidae) og dens parasitoid *Platygaster robiniae* Buhl et Duso, 2007 (Platygastridae) to nye arter for Danmark [*Obolodiplosis robiniae* (Haldeman, 1847) (Cecidomyiidae) and its parasitoid *Platygaster robiniae* Buhl et Duso, 2007 (Platygastridae) two species new for Denmark]. *Entomologiske Meddelelser*, 77(2), 141-144 (in Danish).
- Laguerre, M. & Dauphin, P. (2007). Présence en France d'*Obolodiplosis robiniae* (Haldeman) (Diptera Cecidomyiidae), gallicole sur *Robinia pseudoacacia* L. (Fabaceae) [Presence in France of *Obolodiplosis robiniae* (Haldeman) (Diptera Cecidomyiidae), living in the galls on *Robinia pseudoacacia* L. (Fabaceae)]. *Bulletin de la Société Linnéenne de Bordeaux*, 35(3), 291-296 (in French).
- Li, G., Xu, G., Guo, K., & Du, S. (2014). Mapping the global potential geographical distribution of black locust (*Robinia pseudoacacia* L.) using herbarium data and a maximum entropy model. *Forests*, 5, 2773-2792. <https://doi.org/10.3390/f5112773>.
- Maslyakov, V.Yu. & Izhevsky S.S. (2011). Invazii rastitel'noyadnykh nasekomykh v yevropeyskuyu chast' Rossii [Alien phytophagous insects invasions in the European part of Russia]. Moscow, IGRAS (in Russian).
- Mihajlović, L., Glavendekić, M.M., Jakovljević, I., & Marjanović, S. (2008). *Obolodiplosis robiniae* (Haldeman) (Diptera: Cecidomyiidae) – nova štetočina bagrema u Srbiji [*Obolodiplosis robiniae* (Haldeman) (Diptera: Cecidomyiidae) – a new invasive insect pest on black locust in Serbia]. *Bulletin of the Faculty of Forestry*, 97, 197-207 (in Serbian).
- Mocreaș, S.N. & Timuș, A. (2014). *Obolodiplosis robiniae* (Diptera: Cecidomyiidae) and its parasitoid *Platygaster robiniae* (Hymenoptera: Platygastridae) on *Robinia pseudoacacia* – new invasive species in the Republic of Moldova. In: Sustainable Use and Protection of Animal World Diversity. 176-178.

- Molnár, B., Boddum, T., Szöcs, G., & Hillbur, Y. (2009). Occurrence of two pest gall midges, *Obolodiplosis robiniae* (Haldeman) and *Dasineura gleditschiae* (Osten Sacken) (Diptera: Cecidomyiidae) on ornamental trees in Sweden. *Entomologisk Tidskrift*, 130(2), 113-120.
- Pernek, M. & Matošević, D. (2009). Bagremova muha šiškarica (*Obolodiplosis robiniae*) – novi parazitoida štetnik bagrema i prvi nalaz *Platygaster robiniae* u Hrvatskoj [Black locust gall midge (*Obolodiplosis robiniae*), new pest on black locust trees and first record of parasitoid *Platygaster robiniae* in Croatia]. *Šumarski list*, 3–4(133), 157-163 (in Croatian).
- Petrov, D.L. (2010). Dendrofil'nyye galloobrazuyushchiye dvukrylyye (Insecta: Diptera) fauny Belarusi [Dendrophilous gall midge (Insecta: Diptera) of the fauna of Belarus] *Vestnik Belorusskogo gosudarstvennogo universiteta. Seriya 2. Khimiya. Biologiya. Geografiya*, (1), 31-35 (in Russian).
- Roskam, H., van der Aa, H., Bijkerk, J., Ellis, W., & Moraal, L. (2008). Massaal voorkomen van *Obolodiplosis robiniae* (Diptera: Cecidomyiidae), een nieuwe galmugsoort voor Nederland [Explosive occurrence of *Obolodiplosis robiniae* (Diptera: Cecidomyiidae), a gall midge new for the Netherlands]. *Entomologische Berichten*, 67(1), 27-28 (in Dutch).
- Rupais, A., Stalazs, A., & Strelcuns, R. (2014). Keys to identification of pests by injuries to woody plants in fruit-gardens and parks. *Scripta Letonica (Latvia)*, 1, 5-221.
- Sánchez, I., & Umanan, Á. (2013). Primeros registros de *Obolodiplosis robiniae* (Haldeman, 1847) (Diptera: Cecidomyiidae) en España [First records of *Obolodiplosis robiniae* (Haldeman, 1847) in Spain (Diptera: Cecidomyiidae)]. *BV news Publicaciones Científicas*, 33, 109-111.
- Schneider, N. & Walisch, T. (2009). Sur la présence d'*Obolodiplosis robiniae* (Haldeman, 1847) au Luxembourg (Insecta, Diptera, Cecidomyiidae) [On the presence of *Obolodiplosis robiniae* (Haldeman, 1847) in Luxembourg (Insecta, Diptera, Cecidomyiidae)]. *Bulletin de la Société des naturalistes luxembourgeois*, 110, 161-165 (in French).
- Šefrova, H. & Lantuvka, Z. (2005). Catalogue of alien animal species in the Czech Republic. *Acta universitatis agriculturae et silviculturae mendelianae brunensis*, 53(4), 151-170.
- Sinchuk, O.V., Roginsky, A.S. & Buga, S.V. (2019). Pervaya nahodka *Obolodiplosis robiniae* (Haldeman, 1847) (Diptera: Cecidomyiidae) na territorii Bryanskoy oblasti [The first record of *Obolodiplosis robiniae* (Haldeman, 1847) (Diptera: Cecidomyiidae) from Bryansk area of Russia]. *Euroasian Entomological Journal*, 17(6), 401-402 (in Russian).
- Sitzia, T., Cierjacks, A., De Rigo, D., & Caudullo, G. (2016). *Robinia pseudoacacia* in Europe: distribution, habitat, usage and threats. In: *European Atlas of Forest Tree Species*. Luxembourg, Publication Office of the European Union, 166-167.
- Skrzypczyńska, M. (2007). Muchówki przyszczarkowate (Diptera: Cecidomyiidae) na wybranych stanowiskach w południowej Polsce [Gall midges (Diptera: Cecidomyiidae) in the selected localities in southern Poland. Dipteron]. *Dipteron*, 23, 26-33 (in Polish).
- Skuhravá, M., Bayram, Ş., Çam, H., Tezcan, S., & Can, P. (2005). Gall midges (Diptera: Cecidomyiidae) of Turkey. *Türkiye Entomoloji Dergisi*, 29(1), 17-34.
- Skuhravá, M., Martinez, M., & Roques, A. (2010). Alien terrestrial arthropods of Europe. Chapter 10. Diptera. *BioRisk*, 4(2): 553-602. DOI: 10.3897/biorisk.v4i0.53.
- Skuhravá, M. & Skuhravý, V. (2005). Does the gall midge *Obolodiplosis robiniae* occur in England? *Cecidology*, 20, 34-35.
- Skuhravá, M., Skuhravý, V., & Buhr, H.J. (2013). Gall midges (Diptera: Cecidomyiidae) of Georgia. *Acta Societatis Zoologicae Bohemicae*, 77, 271-309.
- Skuhravá, M., Skuhravý, V., & Csóka, G. (2007). The invasive spread of the gall midge *Obolodiplosis robiniae* in Europe. *Cecidology*, 22(2), 84-90.
- Stalažs, A. (2014). New records of some dipterans (Diptera: Cecidomyiidae, Tephritidae) in north-eastern Lithuania. *Zoology and Ecology*, 24(1), 55-57. <https://doi.org/10.1080/21658005.2014.883831>.
- Tataroğlu, M., Katılmış, Y., & Skuhravá, M. (2024). Three New Records of Gall Midges (Diptera: Cecidomyiidae) from Türkiye. *Journal of the Entomological Research Society*, 26(3), 361–365. <https://doi.org/10.51963/jers.v26i3.2637>.

First Record of the Black Locust Gall Midge Obolodiplosis robiniae

- Tomiczek, C., Cech, T.L., Fürst, A., Hoyer-Tomiczek, U., Krehan, H., Perny, B., & Steyrer, G. (2009). Forstschutzsituation 2008 in Österreich [Forest protection situation 2008 in Austria]. *Forstschutz Aktuell*, 3-8 (in German).
- Tomov, R., Trencheva, K., Trenchev, G., Çota, E., Ramadhi, A., Ivanov, B., Naceski, S., Papazova-Anakieva, I., & Kenis, M. (2009). Non-indigenous insects and their threat to biodiversity and economy in Albania, Bulgaria and Republic of Macedonia. Sofia–Moscow, Bulgaria, PENSOFT. 114 p.
- Tóth, P., Váňová, M., & Lukáš, J. (2009). The distribution of *Obolodiplosis robiniae* on black locust in Slovakia. *Journal of Pest Science*, 82(1), 61-66. <https://doi.org/10.1007/s10340-008-0220-2>.
- Vidano, C. (1970). Foglioline di *Robinia pseudoacacia* con mine di un Microlepidottero nuovo per l'Italia [*Robinia pseudoacacia* leaves with new microlepidopteran mines for Italy]. *L'apicoltore moderno*, 61(10), 1-2 (in Italian).
- Wehrmaker, A. (2007). The black locust gall midge *Obolodiplosis robiniae* (Diptera: Cecidomyiidae) already widespread in southwestern Germany in 2006. *Cecidology*, 22, 57-59.
- Wermelinger, B. & Skuhravá, M. (2007). First records of the gall midge *Obolodiplosis robiniae* (Haldeman) (Diptera: Cecidomyiidae) and its associated parasitoid *Platygaster robiniae* Buhl & Duso (Hymenoptera: Platygastridae) in Switzerland. *Bulletin de la Société Entomologique Suisse*, 80, 217-221.
- Whitebread, S. E. (1989). *Phyllonorycter robinella* (Clemens, 1959) in Europe (Lepidoptera, Gracillariidae). *Nota Lepidopterologica*, 12, 344-353. <https://doi.org/10.2478/v10043-008-0018-8>.
- Zhao, J.-Q., Gao, T., Du, J. J., & Shi, J. (2023). Future Trends in *Obolodiplosis robiniae* Distribution across Eurasian Continent under Global Climate Change. *Insects*, 14, 48. <https://doi.org/10.3390/insects14010048>.