

PEDOLOGICAL EROSIONAL PROCESSES IN REPUBLIC OF MOLDOVA AND THEIR ESTIMATION

Kuharuk E. S.

Institute of Pedology, Agrochemistry and Soils Protection
«N. Dimo», Chisinau

Moldova is the unique world country that possesses the most fertile soils – chernozems that occupy 75 % of its area. But, ill-considered soils' use led to intensive degradation of their ecological functions: energetic, biogeochemical, hydrological, gas-atmospheric, geological etc.

First of all, ecological territorial balance was disturbed. In the whole of the country, natural ecosystems (wood plantations, including forest shelter belts, meadows, grasslands, swamps and water systems) quota makes just 19 %, and at some villages it falls until 7 % of total territory's area. The country's woodiness makes only 9,6 %, and at some regions falls to 1,0 %. At the same time, agricultural lands' cultivation level has increased and rose up to 70 % in average, while in some villages exceeding 90 %.

The actual state of cultivated soils is no less alarming. Negative processes, which lead to soils' degradation and destruction, increase together with exploitation intensification. Soils' structure's damage, increasing compactible degree, dehumification, erosion, hydromorphism, landslide processes, pollution are developing more intensely. Intense cultivation led to ploughing layer's natural structure's destruction, which worsened water and air penetrability along with other physical characteristics of soil. In consequence of applied tractors and other agrotechnique increasing weight a systematic increase of soil's density is observed. Moldova's chernozems and wood soils are shrinking ad litterarum, which is also a negative influence on their characteristics.

Energetic function, reviewed as soils humus potential, plays an important role in soils' ecological functions system, which is important not only for Moldova, but also for the other countries, where other energy sources are limited. In early 60s Moldova's soils contained nearly 1 milliard ton of humus on the whole area in 1 meter layer, now it has 0.8-0.9 milliard ton. Two thirds of this huge loss is ecologically useless, because of erosion.

Water erosion in Moldova's conditions is the most wide-spread soil-destructing process. Averagely and strongly washed-off soils make more than 275 thousands of hectares. It should be noted that the more eroded soils is, the easier it is exposed to further washing off, because lower horizons have less anti-erosion resistance. Erosion processes' intensity has particularly increased in last years as a consequence of wood protection belts' destruction, implementation of so-called intra-farm crops rotation, row crops' quota

increasing in cultivated areas structure and their cultivation by industrial technologies. Applying herbicides to fight weeds excludes cultivation and also excludes possibility to realize appropriate agrotechnical anti-erosion measures: slitting, discrete furrowing, chiseling, and hollowing.

Erosion processes intensity on deep-ploughed soils has increased sharply because transitional chernozems horizons appear on the surface – less humified, with worse structure, water penetrability, and low anti-erosion resistance. Erosion processes are the reason of republic's annual losses of 50 thousands tones of winter grain crops, 100 thousands tones of maize, 25 thousands tones of sunflower seeds, 150 thousands tones of sugar beet.

It was calculated that if we take a full-profiled chernozems potential as 100, than weak wash-off one will have 75, averagely wash-off one – 60–65, strongly wash-off one – 34. Erosion completely disturbs soils' potential energy's accumulation process.

Thanks to their biogeochemical function, chernozems have accumulated in their profile carbon in organic form, nitrogen, phosphorus, some microelements. Every 100 ton erosion caused humus losses mean that also losing 5–6 tons of nitrogen, irretrievably almost half of phosphorus (which is the most scarce biofile in the world) quota, the low Clarke elements (manganese, boron, cuprum, zinc, molybdenum etc). In eroded chernozems these substances' losses make 25–75 % from their initial proportion.

Moldova's soils fertility has decreased in half during the century; consequently cultures yield is decreasing accordingly.

In present for soils fertility restoration agrochemists recommend introducing 10–12 tons of manure in soil per hectare or 18–20 million of tons yearly.

Chernozems hydrological function operates with liquid effluent and solid elution. Liquid effluents may be powerful enough to cause erosion drought, and solid elution sharply disperses soils status as bio-inert carcass body. Soil's biota suffers severely too- the quantity of invertebrate, bacteria, fungi, their activity and specific diversity is reduced in 2–4 times, so on every part of washed-off chernozems the quantity of living beings is reduced, like a fragment of biosphere is dying. The prognosis is quite imposing – all eroded agricultural lands will increase in area, and their growth in last decade makes more than 30 % (table).

Downhill and strongly wash-off lands ploughing is observed in 2001 in Stefan-voda, Anenii Noi and other regions. Chernozems' erosion should be minimized. Agronomic views on soils' erosion must give up to a more biosphere-oriented and ecological view, as 1251,8 thousands of people became proprietors of more than 2 millions of hidelands at the end of 2000. While parting up land into hidelands, anti-erosion requirements of farm's

territory organizing were not met, hidelands are often allotted along the hills, which significantly impedes implementation of anti-erosion measures to maintain soil cover. In sum, erosion not only damages soil, but biosphere in whole, brings on negative social and demographic consequences. Several measures – organization, agrotechnical, hydrotechnical, phytomeliorative and others should be taken in order to prevent Moldova’s soils’ ecological functions degradation. Their implementation should be complete, and not fragmented.

Table

Eroded agricultural lands by zones in Republic of Moldova`on 1.01.20011

Zones	Total area, hectares	Eroded soils:									
		total		weakly-		averagely-		strongly-		averagely-+ strongly -	
		ha	%	ha	%	ha	%	ha	%	ha	%
Northern	875176	296685	34,0	194340	22,2	73515	8,4	28881	3,4	102396	11,8
Central	696345	314748	45,2	165730	23,8	102363	14,7	46655	6,7	149018	21,4
Southern	732411	285640	39,0	157468	21,5	91551	12,5	36621	5,0	128172	17,5
South-Eastern	234753	51880	22,1	36152	15,4	12442	5,3	3287	1,4	15729	6,7
Total	2538685	949468	37,4	553433	21,8	279255	11,0	116780	4,6	396035	15,6
Referring to											
Right bank	2274453	889481	39,1	511238	22,5	265486	11,7	112756	5,0	378242	16,7
Left bank	264232	59987	22,7	42195	16,0	13769	5,2	4024	1,5	17793	6,7

One of the first conditions is to change correlation between natural and anthropogenic, transformed landscapes in different regions. Ecological balance is off the question in conditions of complete lack of forests and natural gramineous vegetation in Belti Steppe and Southern Region. Consequently, there’s a strong need in wood planting, hills tufting, flood plains, hollows, runnels, swamps and others natural destination’s restoration. It is necessary to examine Moldova’s flood plains’ soils and to evaluate them.

The other condition is to create natural carcass, a green skeleton – protection belts situated on watersheds. They would serve to smooth climatic conditions; lessen wind’s force and storms’ destructive power. Anti-erosion, flow–regulating belts are strictly necessary in order to regulate surface flow on hills. These belts should be connected with areas of natural forests and wood plantings, created on ravine’s and landslide’s hills, with areas where natural gramineous vegetation is still preserved, and with protection zones along rivers and lakes. Natural carcass will provide a possibility to cultivate soil and plant cultures across the hills and by horizontals, to implement

agrotechnical and anti-erosion measures. It would benefit to preserve local flora and fauna genetic fund.

The areas with mainly distribution of averagely and strongly washed-off soils should be taken out of ploughing fund and intense use; because they practically cannot be protected from erosion while being a part of crops rotation. These soils should be tufted and planted with multiyear herbs. Natural vegetation restoration in rivers' flood plains, cavins and hollows would regulate flow and clean surface waters. Thus, a considerable balance between natural and anthropogenic components of ecosystems will be created and chernozems' ecological functions will be preserved.

Hills area has been impacted by greatest sedimentation. This regional differentiation is consistent with the decrease in clay and increase in sand content of substratum layers as moving from the North to the South.

Apart from using soil, humankind has no other possibility to obtain food products and numerous materials necessary for existence. Soil as a source of material values, as a priceless natural wealth and the main environmental component must function perfectly and serve forever to us and our future generations.